Recent and possible future trends in abortion

F Lafitte *Department of Social Administration, University of Birmingham*

The regularly published abortion statistics are insufficiently detailed to make it easy to assess the significance of increases or decreases in the annual number of abortions. Numbers of abortions are related only to a few broad age groups of the women at risk, the size of those age groups is not normally stated, and the age and marital status groupings employed differ from those used for comparable data about trends in child bearing.

By matching demographic data on trends in pregnancies which are deliberately terminated (abortion) and those which are not (fertility), this note attempts a) to assess the true impact of abortion among women living in England and Wales during 1973-76, and b) to consider whether the annual volume of abortions may increase over the next decade and what such an increase might signify.

Demographers assess a population's reproductive achievement - 'fertility' - by the annual rate at which women of any age (or age group) give birth to babies. Such age-specific fertility rates, measuring the output of babies per 1000 relevant women, are shown in appended Table II, section (a), for 1973-76 for three broad age groups, with married women distinguished from others.* Reproductive wastage - stillbirths and spontaneous or deliberate abortions - can be assessed in the same way. For the same age groups, section (b) of Table II shows the age-specific abortion rates yielded by the statistics of operations performed under the Abortion Act 1967.

These rates are among the lowest reported from non-Communist countries with comparably liberal abortion laws.¹ They also appear to have reached a peak in 1973 (1974 for the non-married aged 16-19) and to have declined somewhat subsequently, with the sole exception of the small group of teenage wives who recorded a continuing slow rise. With this exception (and with the exception of girls under 16, discussed later), women are tending to resort to abortion less frequently.

It does not, however, follow that *pregnant* women are making diminished use of abortion. Thus, for the whole population at risk (women aged 16-49) column 9 of Table II shows a 10 per cent decrease in the overall abortion rate, from 10.1 abortions per 1000

women in 1973 to 9.1 in 1976; but it also shows a 14 per cent decrease in fertility (from 63.5 to 54.6 births per 1000 women). During the 1970s women have very substantially reduced the rate at which they become pregnant at all; and this change would in itself tend to reduce the abortion rate even if there were no reduction in the proportion of pregnancies terminated by abortion.

Known pregnancy rates

It would therefore be more instructive to relate recorded numbers of abortions, not to all women in the relevant groups, but only to those women experiencing pregnancy. The numbers of pregnancies which result in births are known. So are the numbers of those terminated by lawful abortion (see Table I). Added together, these yield annual totals of known pregnancies, from which can be derived the *known pregnancy rates* in section (c) of Table II. The overall known pregnancy rate, col. 9 shows, was reduced by 13.6 per cent from 1973 to 1976, against only a 10 per cent cut in the abortion rate.

These figures take no account of unknown numbers of other pregnancies which are terminated spontaneously (miscarriages) or by illegal abortion. The latter cannot be numerous, but miscarriages are generally thought to account for at least 15 per cent of all conceptions. In the absence of data about them, it is only possible to relate abortions to known pregnancies, so deriving a set of *abortion incidence rates* - i.e. percentages of *known pregnancies* deliberately terminated. Since the incidence of miscarriage is unlikely to fluctuate widely from year to year, trends in abortion incidence rates restricted to known pregnancies can reasonably be used to assess the true impact of abortion.

Abortion incidence rates

Abortion incidence rates so calculated are shown in section (d) of Table II. They do *not* confirm the view that the impact of abortion is declining. They suggest instead:

a) that in the major groups the rate at which known pregnancies are deliberately terminated is becoming stable, at about 5½ per cent for married women aged 20-34 and 29 per cent for those over 34, and

*The non-married comprise unmarried and formerly married (widowed or divorced) women.

TABLE I *Women at risk, births and abortions*

Thousands										
		16-19		20-34		35-49		16-49		
		M	NM	M	NM	M	NM	M	NM	ALL
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
(a) WOMEN										
Actual numbers:										
1973		148	1185	3788	1170	3771	547	7707	2902	10 609
1974		133	1210	3788	1189	3735	537	7656	2936	10 592
1975		135	1249	3802	1205	3698	541	7635	2995	10 630
1976		132	1284	3823	1249	3649	545	7604	3078	10 682
Projected numbers:										
1981		145	1452	3866	1448	3690	572	7701	3472	11 173
1986		143	1376	3907	1582	4046	640	8096	3598	11 694
(b) LIVE BIRTHS										
Actual numbers:										
1973		52.5	19.0	525.4	33.8	40.0	3.4	617.9	56.2	674.2
1974		47.9	19.3	500.4	32.4	35.2	3.2	583.4	54.9	638.4
1975		43.0	19.0	473.3	31.3	32.2	3.1	548.5	53.4	601.9
1976		38.1	18.6	462.5	31.2	29.9	2.8	530.5	52.6	583.1
Projected numbers:										
1981		46.4	26.2	516.0	40.0	33.9	3.8	596.3	70.0	666.3
1986		50.1	25.0	618.2	48.0	39.7	4.2	708.0	77.2	785.2
(c) ABORTIONS										
Actual numbers:										
1973		0.9	23.0	30.6	34.6	15.3	3.1	46.8	60.7	107.5
1974		0.9	23.7	29.8	34.1	14.4	3.1	45.1	60.9	106.0
1975		0.9	23.7	28.5	32.9	13.6	3.0	43.1	59.6	102.6
1976		0.9	23.2	27.0	31.5	11.9	3.0	39.8	57.7	97.5
Projected numbers:										
1981	A	1.2	29.9	25.9	40.0	13.8	3.9	40.9	73.8	114.7
1986	A	1.3	30.9	31.0	48.0	16.2	4.5	48.5	83.4	131.9
1981	B	1.3	34.9	27.0	41.7	16.7	4.5	45.0	81.1	126.1
1986	B	1.4	36.1	33.2	49.1	19.6	5.1	54.2	90.3	144.5

Notes—Births and abortions occurring in England and Wales to women resident there. M: currently married women. NM: non-married women (= all others).

at about 55-56 per cent and 50 per cent respectively for the non-married aged 16-19 and 20-34;

b) that in two much smaller groups incidence rates are still rising: teenage married girls (from 1.7 to 2.3 per cent over 1973-76) and non-married women over 34 (from 47.7 to 51.7 per cent).

The age-specific and incidence rates for abortion during 1973-76 in Table II can be used for projecting possible future trends. If nothing else were to change, then over the decade 1976-86 changes in the size and composition of the relevant age groups could by themselves generate an increase in abortions. Section (a) of Table I shows expected numbers of women in the various age groups in 1981 and 1986, derived from the most recent official population projections (OPCS²). By 1986 the total female population at risk will have grown by 1m., or 9½ per cent; but in the most abortion-prone group, non-married women of 20-34 (and

predominantly under 25), the increase is expected to be disproportionately larger: almost 27 per cent. If in 1986 age-specific abortion rates were to remain unchanged at the 1976 levels (cf. section (b) of Table II), there would nevertheless be 110 000 abortions as against the 97 500 recorded for 1976. If nothing else changes, a 13 per cent increase in abortions can be expected by 1986 simply in consequence of age-group changes; and this increase would reflect no change at all in the true impact of abortion.

This estimate is, however, too simple. It assumes that pregnancy rates as well as abortion rates would remain at 1976 levels. Future pregnancy rates are unpredictable. But there are good reasons for expecting fertility – and therefore pregnancy – rates soon to turn up from the low point they reached in 1976.

The official population projections incorporate

TABLE II* *Pregnancy, births and abortion rates*

	16-19		20-34		35-49			16-49	
	M	NM	M	NM	M	NM	M	NM	ALL
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
(a) BIRTHS PER 1000 WOMEN									
Actual									
1973	355	16.0	139	28.9	10.6	6.2	80.2	19.4	63.5
1974	360	16.0	132	27.2	9.4	6.0	76.2	18.7	60.3
1975	318	15.2	125	26.0	8.7	5.7	71.8	17.8	56.6
1976	287	14.5	121	25.0	8.2	5.1	69.8	17.1	54.6
Projected									
1981	320	18.0	133	27.6	9.2	6.6	77.4	20.2	59.6
1986	350	18.2	158	30.3	9.8	6.6	87.5	21.5	67.1
(b) ABORTIONS PER 1000 WOMEN									
Actual									
1973	6.1	19.4	8.1	29.6	4.1	5.7	6.1	20.9	10.1
1974	6.7	19.5	7.9	28.7	3.9	5.8	5.9	20.7	10.0
1975	6.7	19.0	7.5	27.3	3.7	5.5	5.6	19.9	9.7
1976	6.8	18.1	7.0	25.2	3.3	5.5	5.2	18.7	9.1
Projected									
1981 A	8.3	20.6	6.7	27.6	3.7	6.8	5.3	21.3	10.3
1986 A	9.1	22.5	7.9	30.3	4.0	7.0	6.0	23.2	11.3
1981 B	9.0	24.0	7.0	28.8	4.5	7.9	5.8	23.4	11.3
1986 B	9.8	26.2	8.5	31.0	4.8	8.0	6.7	25.1	12.4
(c) KNOWN PREGNANCIES PER 1000 WOMEN									
Actual									
1973	361	35.4	147	58.5	14.7	11.9	86.2	40.3	73.7
1974	367	35.5	140	55.9	13.3	11.8	82.1	39.4	70.3
1975	325	34.2	132	53.3	12.4	11.2	77.5	37.7	66.3
1976	295	32.6	128	50.2	11.5	10.6	75.0	35.8	63.7
Projected									
1981 A	328	38.6	140	55.2	12.9	13.5	82.7	41.4	69.9
1986 A	359	40.6	166	60.7	13.8	13.6	93.4	44.6	78.4
1981 B	329	42.0	140	56.4	13.7	14.5	83.3	43.5	70.9
1986 B	360	44.4	167	61.4	14.7	14.5	94.1	46.6	79.5
(d) ABORTIONS % OF KNOWN PREGNANCIES									
Actual									
1973	1.7	54.8	5.5	50.6	23.4	47.7	7.0	51.9	13.8
1974	1.8	55.1	5.6	51.3	29.0	49.2	7.2	52.6	14.2
1975	2.1	55.5	5.7	51.2	29.7	49.2	7.3	52.7	14.6
1976	2.3	55.5	5.5	50.2	28.5	51.7	7.0	52.3	14.3
Projected									
1981 A	2.5	53.3	4.8	50.0	29.0	50.6	6.4	51.3	14.7
1986 A	2.5	55.3	4.8	50.0	29.0	51.7	6.4	51.9	14.4
1981 B	2.7	57.1	5.0	51.0	33.0	54.2	7.0	53.7	15.9
1986 B	2.7	59.1	5.1	50.6	33.0	54.8	7.1	53.9	15.5

*A thoroughly refined analysis would establish the number of known pregnancies more precisely a) by counting stillbirths as well as live births, and b) by adding to the births and stillbirths occurring in a year the abortions occurring, not in the same period, but six months earlier (ie resulting from the same 'cohort' of pregnancies). The effect of (a) is negligible. Neglect of (b) - inevitable without unpublished data - is unlikely to distort the picture of long-term trends.

the assumption that fertility will not remain static but will rise until by the mid-1980s it reaches replacement level - i.e. suffices to yield completed families averaging 2.1 children per woman (as

against about 1.7 for 1976 fertility). 'A simple assumption of replacement-level fertility may be regarded as the most appropriate basis for a long-term projection, particularly when the available

evidence about the intentions of young married women at the present time points to average family size somewhere in that neighbourhood' (OPCS²). The projected births in Table I, section (b), and the fertility rates in Table II, section (a), are derived from the trends assumed in the official projections.

The hypotheses

If fertility moves in this way and the pattern of births is as shown, it becomes possible to estimate future trends in known pregnancies by applying differing assumptions about abortion incidence rates. The two hypotheses explored in this note are that abortion incidence rates a) remain stable at about the 1976 level or b) increase appreciably. The key assumptions are:

		<i>Abortions % of known pregnancies</i>		
		<i>1976 actual</i>	<i>A</i>	<i>B</i>
All women:	16-19	29.8	30.0	33.3
	20-34	10.6	10.6	11.0
	35-49	31.3	32.0	36.0
Non-married women:	20-34	50.2	50.0	51.0

Given these parameters, rates for all other groups were worked out by a process of 'best fitting'.

The outcome, in section (c) of Table A, is striking. On hypothesis *A*, assuming 1976 fertility as well as no change in the true impact of abortion, age-group effects alone would raise the number of abortions by 13 per cent to 110,000 by 1986. If, however, fertility were to rise moderately up to replacement level,* then the abortion total would grow to 132 000 – an increase of 35 per cent (matching a 35 per cent increase in births).

Thus, over the next ten years changes in fertility and pregnancy rates may be expected to influence the number of abortions much more than changes in the size of relevant female age groups. And an increase in abortions among women over 15 from 97 500 a year to anywhere between 110 000 and 132 000 could be entirely compatible with no increase at all in the true impact of abortion, in terms of the proportions of pregnancies terminated.

Hypothesis *B* allows for some increase in the incidence of abortions in all groups, up to the levels shown for 1986 in the last line of section (d) of

*Such an increase would be moderate. Fertility measured in terms of the implied average completed family size ('total period fertility rate') rose from 2.14 in 1951 to 2.94 in 1964, then dropped to 2.02 in 1973 and 1.72 in 1976. The corresponding net reproduction rates were 1.00 (= exact generation replacement), 1.37, 0.96 and about 0.82 (OPCS³).

Table B – for example from 55½ to 59 per cent among non-married girls of 16-19 and from 28½ to 33 per cent among married women over 34. On this hypothesis the number of abortions could increase by nearly 50 per cent to 145 000.

On both hypotheses the established tendency for the non-married to obtain more abortions than the married would be intensified. The non-married share of abortions rose from 57 per cent in 1973 to 59 per cent in 1976, and may rise to 63 per cent in 1986, with girls of 16-19 accounting for 23-25 per cent of the total, against 21 per cent in 1973. This shift would come about chiefly because, as already noted, the most abortion-prone group, non-married women aged 20-34 (with a pregnancy rate of 50 per 1000 women, with half the pregnancies being terminated), is also the group which will experience the most disproportionate numerical enlargement.

How plausible are these hypotheses? It is not implausible to expect some increased resort to abortion among unmarried girls in their 'teens, and possibly also among teenage wives – all beginners in adult life, sex and the arts of contraception. It may also be plausible to expect some increase among the non-married over 34. To that extent – but no further – hypothesis *B* may be justified. But for all the other groups there are no current indications of any increase at all, and a good many indications that reduced rather than increased resort to abortion is more likely. Thus it is highly likely that couples who have completed their wanted families will turn increasingly to sterilization of husband or wife so that abortion rates and incidence among married women over 30 or 35 are more likely to fall than to remain stable (hypothesis *A*) or to rise (hypothesis *B*).* The incidence of abortion among all adult age groups, and eventually perhaps also among teenagers, is also more likely to fall as knowledge and practice of reliable contraception become more firmly established, now that the National Health Service has assumed full responsibility for family planning services, for the unmarried as well as the married, and over 90 per cent of general practitioners are undertaking to give advice.

It may be fairly concluded that hypothesis *B* is not realistic, and that the hypothesis *A* estimates based on continuance of the existing incidence of abortion, is more likely to overstate than to understate future trends. They are best regarded as maxima not likely to be attained.

Under age pregnancies

Excluded from the foregoing review are girls under 16. The gist of the little information published about them is summarised on the next page.

*Only one fifth of children are now born to women aged 30 and over, against three fifths in mid-Victorian times.

	Ages	Rates per 1000 girls		Abortions	
		Births	Abortions	Known pregnancies	% of known pregnancies
1973	12-14	0.29	0.75	1.04	72
	15	3.75	6.39	10.14	63
	12-15	1.16	2.16	3.32	65
1975	12-15	1.00	2.41	3.41	71
1976	12-15	0.79	2.24	3.03	74

In this school-girl group the bulk of births and abortions occur among 15-year-olds and most of the rest among 14-year-olds. Even among girls of 15 pregnancies are, fortunately, rare: about 99 per cent do *not* become pregnant at that age; and it is not clear whether there is any tendency for the incidence of pregnancy to increase.

What is clear, taking the age group 12-15 as a whole, is that when pregnancy does occur it is now normally and increasingly ended by abortion. The abortion incidence rate rose from 65 per cent in 1973 to 74 per cent in 1976. This is a trend which may be expected to progress further, since it appears to be favoured by the climate of opinion. By 1986 at least 80 per cent, and possibly 90 per cent, of these premature pregnancies may be terminated.

In 1976 three girls per 1000 in the 12-15 age group were pregnant and 74 per cent of their pregnancies were terminated. The outcome was 3500 abortions and 1200 births. If in 1986 the pregnancy rate is the same, the outcome will be:

a) if 80 per cent are terminated: 3250 abortions and 800 births

b) if 90 per cent are terminated: 3650 abortions and 400 births.

Although there are no current indications that this may happen, if the pregnancy rate were to rise to 3.5 per 1000 (it did reach 3.4 in 1975), and if the termination rate rose to 90 per cent, then in 1986 there could be 4250 abortions and 470 births occurring among girls aged 12-15.

References

- ¹Tietze, C and Murstein, M C (1975). *Induced Abortion: 1975 Factbook*, Reports on Population/Family Planning, no. 14, 2nd ed., Population Council, New York.
- ²Office of Population Censuses and Surveys (1977). *Population Projections 1975-2015*, Series PP2, no. 7, HMSO, London.
- ³Office of Population Censuses and Surveys (1977). *Birth Statistics 1974, England and Wales*, Series FM1, no. 1, HMSO, London, Table 1.4.