The assessment of pain in advanced cancer

Robert G Twycross *The Churchill Hospital, Headington, Oxford*

This is one of a group of papers read at the London Medical Group conference on 'Pain: a necessity?', which was held in Charing Cross Hospital Medical School, London in February 1978. Dr Twycross argues that complete assessment implies the ability not only to make a diagnosis but also to initiate appropriate treatment. Describing the site, severity and quality of the pain is only the first step. A doctor needs to:

- 1) *Be aware of the range of diagnostic possibilities*
- 2) *Appreciate the influences of non-physical factors such as mood and morale*
- 3) *Be aware of the range of treatment options*
- 4) *Establish realistic objectives with the patient*
- 5) *Reassess at appropriate intervals to review treatment, monitor side-effects, and develop and maintain a good working relationship with the patient.*

Traditionally we are taught to assess pain by determining its PQRST characteristics (Table 1). Unfortunately a blind belief in the efficacy of so simple an approach may hinder rather than help the doctor in the assessment of the patient's pain. Determination of the PQRST characteristics is only the beginning, providing a description of the pain but no more. Complete assessment implies the ability to make a diagnosis. This demands a grasp of both general and neurological anatomy, an understanding of the phenomenon of referred pain and a knowledge of the range of pathological processes which are potential causes of pain. From the history and clinical examination, supplemented if necessary by x-ray scan, or other test, it should be possible to develop a fairly clear mental picture of the physical mechanisms underlying the pain. Then, assessment completed and diagnosis made, treatment is initiated. We are familiar with this

sequence of events in relation to, say, acute abdominal pain, but often fail to apply the same relentless logic in assessing pain in other circumstances.

My own clinical involvement is largely confined to advanced cancer. Assessment needs to be as thorough in this area as in any other. Cancer can cause pain in any part of the body and by a variety of mechanisms. Many patients have more than one pain. Moreover, alleviation of one pain may unmask another or a new pain may develop. To cope with what is often a complex situation, I find the use of a body chart to record pain data a great help (Fig. 1). Generally, patients put on a brave face for the doctor. Patients in severe pain do not always look distressed. Accordingly, intensity of pain is assessed not only by the patient's description but also by discovering what drugs have failed to relieve, whether sleep is disturbed, and in what way activity is limited ("How long is it since you went out?", "What are you doing around the house?"), etc. In addition, the patient's spouse should be interviewed. Often it is only the latter's comments that give the true picture - though, when the pain is relieved, the patient frequently concurs spontaneously with the spouse's earlier opinion. A patient who is obviously in distress and who says only "It's all pain, doctor" is best thought of as having 'overwhelming' pain, that is, severe pain compounded by anxiety, depression and loss of morale.

Diagnostic possibilities

A diagnosis of cancer does not necessarily mean that the malignant process is the cause of the pain. A peptic ulcer, cystitis or a musculo-skeletal disorder may be responsible and are all conditions that benefit from more specific treatment. If the pain is due to the cancer, it is important to determine the mechanism(s) of the pain(s) as treatment frequently varies according to mechanism.

Cancer pain may be caused by soft tissue infiltration, bone involvement, nerve compression, elevated intra-cranial pressure, lymphoedema and visceral distension or obstruction. Infection in a malignant ulcer and muscle spasm secondary to bone involvement or deformity also occur. Constipation and bedsores are common indirect forms of cancer pain. Asking the patient to point to the site of the pain and to describe it will usually yield sufficient clues to make the correct diagnosis.

Table 1 *The PQRST characteristics of pain*¹

'Tell me about your pain.' 'Where is it?'		
P	alliative factors	'What makes it less intense?'
Q	provocative factors	'What makes it worse?'
R	quality	'What's it like?'
S	radiation	'Does it spread anywhere else?'
T	severity	'How severe is it?'
	temporal factors	'Is it there all the time, or does it come and go?'

<p style="text-align: center;">BODY CHART</p> <p>Ca. Prostate</p>	<p>Hospital No.</p> <p>Surname</p> <p>First Names</p> <p>R.M. age. 65 13.7.77</p>
--	--

Path ✕ L. femur 26/5, R. humerus 30/5, L. humerus 20/6 → all pinned etc.

Pains developed over last 2/12

Lies flat in bed.

Every movement hurts.

Morphine gives relief 2+ hours.

1. bedsore

2. pressure on heels

3. adductor spasm

4. L. ulna

5. R. radius

6. } ✕ site

7. } ✕ site

Note jumps when touched. Use analgesics + diazepam.

Fig 1 Body chart used to record pain data relating to 65-year-old man with cancer of the prostate gland.

A four-year-old child with an inoperable pontine glioma experienced increasing pain in the head and occipital region. She lay flat all the time because elevation of the head caused a marked increase in pain. With this history it was necessary to postulate a local source of pain (possibly caused by post-radiation meningeal adhesions) in addition to the diffuse headache of secondary hydrocephalus (which would have been helped by a more erect posture). The diffuse pain was relieved by small regular doses of morphine but not until she was transferred from a King's Fund to an Ellison bed (which elevates head, neck and trunk in unison) was it possible for the child to sit up without pain. Subsequently it became possible to transfer the child from bed to a high-backed reclining chair and, eventually, to lift her onto her mother's lap. This suggested that some of the pain had been caused by spasm of the neck muscles, and that the confidence engendered by the ability to sit up in bed allowed additional manoeuvres to be undertaken without pain.

Non-physical factors

Pain is a dual phenomenon, one part being the perception of the sensation and the other the patient's psychological reaction to it. It follows that a person's pain threshold will vary according to mood and morale (Fig. 2). Attention must therefore be paid to factors that modulate pain threshold, such as anxiety, depression, and fatigue (Table II). Much can be done to alleviate pain by explaining the mechanism underlying the pain (this reduces

Fig 2 For any given noxious stimulus the pain experienced varies from ache to agony and depends on the psychological reaction of the sufferer to his discomfort.

Table II Factors affecting pain threshold

discomfort		relief of symptoms	
insomnia		sleep	
fatigue		rest	
anxiety		sympathy	
fear		understanding	
anger		diversion	
sadness		elevation of mood	
depression			
mental isolation			
introversion (past experience)	threshold lowered	analgesics tranquillisers antidepressives	threshold raised

anxiety) and by a continuing concern for the patient (this raises morale). Ignoring mental and social factors may result in otherwise relievable pain remaining intractable. At a weekly seminar for trainee practitioners² discussion once centred on the problem of a patient with metastases in bone from disseminated breast cancer whose pain was not relieved by narcotic analgesics. During the seminar it was suggested that the pain was intractable as the woman was angry because her doctors and relatives would not admit that she was dying nor discuss the problems this created. This proved to be the right explanation; a full and frank discussion resulted in a dramatic improvement in her mental state and she no longer complained of pain.

Treatment options

Relief of pain may be achieved by one or more of the following methods:

- 1) Modification of the pathological process
- 2) Elevation of pain threshold
- 3) Interruption of pain pathways
- 4) Immobilisation

Even in far-advanced cancer, modification of the pathological process by means of radiation, chemotherapy, or hormone treatment should always be considered. As in other areas of oncology, it is important to ensure that the treatment is not worse than the disease. If radiation or an oestrogen is prescribed, this does not mean that analgesics should be withheld. The best results are obtained by adopting a 'broad-spectrum' approach, using two or more treatments in combination (Table III).

The use of analgesics and other drugs is best seen as but one way – generally a powerful way – of elevating the patient's pain threshold. Their use is fully discussed elsewhere.^{3,4} Interruption of pain pathways refers to chemical neurolysis (nerve blocks) and neurosurgical techniques (e.g. rhizotomy, spinothalamic tractotomy). Such procedures are of value in nerve compression/infiltration pain that is inadequately relieved by the use of analgesics and corticosteroid.

Some patients continue to experience pain on movement despite analgesics, other drugs, radiotherapy and nerve-blocks. In these, the situation may be improved by suggesting commonsense modifications to daily activity. For example, a man may continue to struggle to stand while shaving unless the doctor suggests that sitting would be a good idea. Such a suggestion is accepted more readily if accompanied by a simple explanation of why weight-bearing precipitates or exacerbates the pain. Individually designed plaster or plastic supports for patients with multiple collapsed vertebrae or Thomas splints for femoral pain are occasionally necessary to overcome intolerable pain on movement in bedfast patients.

Table III Treatment of pain in advanced cancer

Mechanism	Analgesic and adjuvants	Co-analgesic	Non-drug treatment	Other measures
1. Soft tissue infiltration 2. Bone involvement 3. Nerve compression 4. Raised cerebral pressure 5. Lymphodema 6. Abdominal visceral a) epigastric b) hypogastric 7. Ulceration – infection	<div><div><div>morphine (severe)</div><div>codeine (moderate)</div><div>aspirin (mild)</div></div><div>antiemetic ? anxiolytic ? laxative</div></div>	<div><div>aspirin prednisolone dexamethasone diuretic</div><div>antibiotic</div></div>	<div><div>radiotherapy nerve block compression sleeve coeliac axis ganglion block (presacral block)*</div></div>	<div><div>modification of lifestyle</div><div>diversion – heat – massage</div></div>
8. Constipation	specific treatment			
9. Second pathology				

the next day indicated that, although less severe, he was still in pain. Further questioning revealed that the site of pain was now retrosternal and epigastric; he had no femoral pain at all. The dose of morphine was left unaltered, and the prescription of an antacid resulted in complete relief.

In the case of the man with overwhelming pain (Fig. 1), it was necessary to review progress in relation to each pain. On the second day he was reluctant to admit that several of the pains were less intense but, judging by his reactions to passive movement, they undoubtedly were. Possibly his reluctance was due to a shifting base-line of reference. It was necessary to remind him of the difference I observed and to encourage him to accept that the pains were beginning to ease. It was a case of 'chipping away' at his total pain experience until eventually, after a week or so, he was considerably more comfortable and able to sit up in bed.

Acknowledgements

Tables II and III and Fig. 1 are reproduced by permission of Pitman Medical, Tunbridge Wells, England.

References

- ¹Gray, J (1977). A pain in the neck - and shoulder. *Pain Topics*, 1, No. 3, 6.
- ²Anonymous, (1971). Vocational training for general practice. *British Medical Journal*, 2, 704-706.
- ³Vere, D W (1978). Pharmacology of morphine drugs used in terminal care. In: *Topics in Therapeutics* 4, edited by D W Vere, pp. 75-83. Pitman Medical, Tunbridge Wells.
- ⁴Twycross, R G (1978). Bone Pain in Advanced Cancer, *ibid.*, pp. 94-110.
- ⁵Twycross, R G (1977). Care of the terminal patient, In: *Breast Cancer Management - early and late*. Edited by B A Stoll, pp. 157-163. Heinemann Medical Books Ltd, London.