

In defence of genital autonomy for children

Brian D Earp

ABSTRACT

Arora and Jacobs (2016) assume that liberal societies should tolerate non-therapeutic infant male circumcision, and argue that it follows from this that they should similarly tolerate—or even encourage—what the authors regard as ‘*de minimis*’ forms of female genital mutilation (as defined by the World Health Organization). In this commentary, I argue that many serious problems would be likely to follow from a policy of increased tolerance for female genital mutilation, and that it may therefore be time to consider a less tolerant attitude toward non-therapeutic infant male circumcision. Ultimately, I suggest that children of whatever sex or gender should be free from having healthy parts of their most intimate sexual organs either damaged or removed, before they can understand what is at stake in such an intervention and agree to it themselves.

INTRODUCTION

In their target article, Arora and Jacobs¹ contend, among other things, that if non-therapeutic infant male circumcision should be widely tolerated, then so should some forms of what the WHO calls ‘female genital mutilation’ (or FGM).² Indeed, there are numerous substantial overlaps, both physical and symbolic, between these two types of genital alteration that are not very widely appreciated;^{3–10} accordingly, I have prepared an online supplementary appendixⁱⁱ in which I outline some of the main morally relevant features that they share. For now, however, let us consider only the conditional argument put forward by the authors, namely that: ‘a liberal society that tolerates expression of culture and/or religion in the manner of male circumcision should also permit certain *de minimis* [female genital altering] procedures’.¹

I offer a different perspective. Rather than proceeding from the premise that non-therapeutic, non-consensual male genital alteration (MGA) is clearly

permissible (and should therefore be widely tolerated in liberal societies) to the conclusion that purportedly ‘*de minimis*’ forms of non-therapeutic, non-consensual female genital alteration (FGA) should also be permitted, I will instead move in the opposite direction. My strategy will be to highlight some of the downsides that seem likely to follow from an increased tolerance for FGA in Western societies, and use this as a premise from which to argue that it may be time to consider a less tolerant position towards MGA. Please note that I will not be arguing in favour of a ‘ban’ on MGA (for reasons I go into elsewhere)^{11–13} but rather for its being strongly discouraged, perhaps through some form of regulation (akin to the proposal by Ben-Yami in an earlier issue of this journal).¹⁴

SOME PROBLEMS WITH INCREASING TOLERANCE FOR FGA IN WESTERN SOCIETIES

Here are some problems that seem likely to follow from a more tolerant position towards FGA.

Legal problems

FGA, including its ‘minor’ forms, is currently illegal in the vast majority of liberal societies (I will focus on so-called Western countries, such as the USA, Canada, Australia and the countries of Europe).^{15–16} To change this situation, at least according to one plausible interpretation, it would require that the laws regarding physical assault on a minor be rewritten, potentially creating widespread disturbances and inconsistencies throughout their respective legal systems. This is because, as some scholars have argued, cutting into a child’s genitals without a

medical diagnosis, and without its informed consent, meets the formal definition of criminal assault under the legal codes of most of these societies.^{4 17–20} (Please note that this view assumes that parental ‘proxy’ consent is invalid for such procedures, for which arguments have been given elsewhere.)^{20–22} Indeed, in its move to ban FGA in 1997, the US Congress stated ‘that it ‘infringes upon the guarantees of rights secured by Federal and State law, both statutory and constitutional’. That is to say, [non-therapeutic] female genital cutting [without consent] was *already unlawful*’ prior to the enactment of specific legislation.²³

A further issue, as Blackstone noted more than 200 years ago, is that ‘the law cannot draw [a] line between different degrees of violence’²⁴ [p. 270]. What this means is that if any particular act of violence—such as the cutting of a child’s body in the absence of medical necessity, perhaps especially with the effect of damaging or removing functional tissue—is legally permitted, then fraught decisions have to be made ‘as to where within the continuum of violence the limits are to be fixed’²⁵ [p. 129]. This, in turn, could open the door for interested parties to argue, for any proposed act of cutting, that it is not ‘violent enough’ to warrant the placement of limitations on parental decision making.⁴ More on this will be said in a subsequent section.

An additional problem is that changing the law would remove at least one important tool that reformers from within the affected communities can appeal to, in their quest to end the spectrum of FGA practices. Specifically, many women from established immigrant groups report that they do not wish to continue subjecting their daughters to FGA (for an excellent review of the evidence concerning ‘cultural change after migration’ with respect to this issue, see the work of Johnsdotter and Essén)²⁶ but they may face pressure from fellow group members.²⁷ This can make it difficult for any one parent or family to unilaterally challenge, much less abandon, the tradition: as Mackie has argued, it is a collective action problem.^{28–29} What a legal prohibition allows, then, is for reformers from within these groups to be able to point to the law and say, ‘I have no choice but to keep my daughter’s genitals intact’, effectively resolving the social dilemma.^{27–30}

Finally, as Ludbrook argues, ‘the law is an important symbol and if [it] allows parents, carers and teachers to [so much as hit their] children’, much less cut into a physically and symbolically sensitive part

ⁱThe terminology here is notoriously contentious: see, for example, Davis³ for an in-depth discussion. I will simply follow Arora and Jacobs in using ‘female genital alteration’ (FGA) for the remainder of this essay.

ⁱⁱAvailable at <http://dx.doi.org/10.1136/medethics-2015-103030>. Please note that this appendix also includes a more detailed response to many of Arora and Jacobs’s more problematic claims, focusing on those that I did not have the room to address in the main text of this commentary.

of their bodies, 'it is sending a message to these people and to the community generally that children are not entitled to the same right to bodily integrity as adults'²⁵ [p. 129].

Against this view, Arora and Jacobs argue that (what they regard as) '*de minimis*' forms of FGA 'do not constitute a human rights violation' with respect to bodily integrity, because they have 'no more effect than other accepted procedures performed on minors for esthetic enhancement'.¹ In previous writings, they have listed mole removal and cosmetic orthodontia as examples of such procedures.³¹ But as I have suggested elsewhere:

'the genitals (in particular) might plausibly be seen as having a special, even unique psychosexual significance compared to other parts of the body, which could make their un-consented alteration more likely to be experienced (later on) as a harm. [This] could help to explain why there is an active 'genital autonomy' movement in the United States, Europe, and elsewhere—fueled by women, men, and intersex people who are extremely resentful about their childhood genital surgeries—but not an anti-orthodontics movement or an anti-mole removal movement'³² [p. 45].

More could be said about this disagreement. For a thoughtful analysis of the right to bodily integrity as it applies to preautonomous individuals, see the essay by Ungar-Sargon in a recent issue of this journal.³³ See also the work of Darby on a child's 'right to an open future'.³⁴ For an extended articulation of my own views concerning the ethics of ostensible 'enhancement' procedures in children, see the references here.^{5 12 35 36} iii

Regulatory problems

Another problem with the authors' proposal has to do with monitoring and regulation. As Arora and Jacobs concede, their suggestion might 'open the door' for 'more invasive procedures [to be carried out] under the guise of *de minimis* procedures',¹ thereby increasing the overall level of harm. In response to a similar policy proposal by the American Academy of Pediatrics (AAP) in 2010 (see later discussion), the Somali-born writer and

activist Hirsi Ali asked: 'how could we tell that parents who want to ensure that their daughter will be a virgin on her wedding night will not have her (legally) nicked [on one day] and then a few months later (illegally) infibulated?'³⁷ Indeed, as Leye and colleagues point out, 'in some communities, traditional circumcisers will 'redo' girls if they notice that only an incision has been made'³⁸ [p. 368].

The authors' response to this type of objection is that 'given the widespread nature currently' of the more invasive procedures, 'if even a few girls undergo a *de minimis* procedure instead of a more invasive one due to this strategy then the strategy is appropriate'.¹ However, they do not provide any evidence that the more invasive procedures are in fact 'widespread' in Western countries, specifically, which is the legal setting most relevant to their arguments (and readership); in fact the recent survey by Johnsdotter and Essén stands in direct contrast with this scenario.²⁶ Moreover, if permitting 'minor' forms of FGA actually had the effect of lending a 'cloak of respectability' to all FGA procedures—a possibility that Arora and Jacobs themselves raise, but do not adequately consider—then the more invasive forms might very well increase in number, leading (again) to greater harm overall. Since the authors do not provide any persuasive evidence in favour of their own predicted outcome as opposed to this plausible alternative, it is unclear why we should endorse their policy proposal, even on consequentialist grounds.

A related problem is more practical in nature. How would the actual cutting sessions be monitored (and by whom) to make sure that only the 'right amount' of tissue was being incised, damaged, or removed? Arora and Jacobs admit that this is a problem with their proposal. But instead of addressing it, they simply divert attention to other problematic practices. As they write: 'The concern regarding amount of tissue being removed is ... not unique to FGA, but is similar to male circumcision as well as cosmetic surgeries on adults'.¹

They are right to raise the example of male circumcision. At least one underdocumented risk of this procedure, especially when carried out in infancy, is the removal of too much tissue—sometimes causing painful erections when the child grows up.^{39–42} To put it simply, there is no 'dotted line' showing where to cut around an infant's diminutive penis, just as there is no determinate location where the foreskin ends and where the rest of

the penis begins. This uncertainty in terms of where to cut, and the associated risk of cutting away too much, arguably speaks in favour of deferring the surgery until the organ has reached its full size. Then, the individual whose genital integrity is at stake can indicate exactly how much tissue - if any -^{iv} he would like to have removed, and he can do so under conditions of informed consent.

A similar analysis applies to FGA procedures. In fact, this parallel leads us directly to the other diversion mentioned by the authors, namely, 'cosmetic surgeries on adults'. The key word is 'adults'. If an *adult female* wishes to undergo a genital surgery (to choose the most pertinent example) for cosmetic, cultural, religious or other reasons, there is indeed, as Arora and Jacobs point out, a non-trivial risk that too much tissue will be removed, or that the surgery will yield unsatisfactory results in any number of other ways. But this would be mitigated by the fact that the surgery was undertaken voluntarily. In other words, the primary (moral) difference between consensual genital surgery, and both FGA and MGA carried out on children, is that an adult can assess the relevant risks in light of her own goals and values, and determine whether they are 'worth it' to her, all things considered.^{35 43} A child has no such recourse.

Speaking generally, then, the suggestion of harm tolerance when it comes to interventions into children's bodies is untenable. In part, this is due to 'the difficulty it presents in terms of specifying an appropriate threshold for harm that could be measured in an objective way'⁴⁴ [p. 98]. As Van Howe notes, the meaning of 'harm', in practice, therefore, will inevitably be left up to the interpretational vagaries of each provider, perhaps in consultation with the child's parents.⁴⁴ This vagueness creates a problem: 'there are practitioners, especially in cultures where female circumcision is common, who fervently believe that more invasive forms of female circumcision'—that is, forms that even Arora and Jacobs reject—'do not pose risks of physical or psychological harm'⁴⁴ [p. 167]. In fact, these providers could easily make selective appeals to studies from the medical literature that appear to show health-based benefits for FGA,⁴ as well as an absence of serious harms,⁴⁵ much as some supporters of MGA are prone to do.^{46 47} When the

^{iv}The overwhelming majority of men who grow up with intact genitalia do not go on to pursue circumcision later.

ⁱⁱⁱOf particular relevance is the essay, 'Between Moral Relativism and Moral Hypocrisy: Reframing the Debate on 'FGM'', in press at the *Kennedy Institute of Ethics Journal*. The accepted manuscript is available online ahead of print here: https://www.academia.edu/10197867/Between_moral_relativism_and_moral_hypocrisy_Reframing_the_debate_on_FGM_.

many purported cultural benefits of FGA are factored in, Van Howe continues, ‘practitioners could easily convince themselves that any harm is more than offset by the many perceived benefits’⁴⁴ [p. 167].

Medical problems

Potential problems concerning clinical and surgical matters have been raised by Leye and colleagues. They write: ‘it is difficult to avoid damaging the clitoris when performing a [*de minimis*] incision, especially in genitalia that are not fully developed’. Moreover, ‘complications (such as shock, infections, sepsis, and bleeding) are difficult to avoid [entirely], even with [such a minor] incision’³⁸ [p. 368]. Thus, as Goldman has stated, the assumption that ‘*de minimis*’ interventions into a child’s vulva really are essentially harmless is ‘debatable’. As he goes on to write: even if it could be shown that the harm is, in some sense, ‘relatively minor, it is still harm. Certainly, there is physical harm or risk of physical harm any time a cutting instrument contacts sensitive genital tissue’.⁴⁸ He concludes, then, that the ‘burden of proof’ is on those who argue that the risk of harm is trivial.

How Arora and Jacobs could meet this burden is unclear. First, as they note, it is currently illegal in Western societies to perform FGA procedures in children, so it is impossible to collect the very data that would be needed to make their case. But let us just assume that the ban were lifted. How long would it be reasonable to allow for such procedures to be performed in the absence of high-quality long-term follow-up data (concerning such vital issues as the effects of these procedures on, eg, sexual function, sensation and satisfaction) before it became possible to show that they had, in fact, been more harmful than Arora and Jacobs guessed that they would be?

An example should elucidate the danger. Consider ‘procedures resembling elective labiaplasty as performed in Western nations’, which Arora and Jacobs use to illustrate category 2 of their proposed typology. This refers to interventions that ‘create morphological changes, but are not expected to have an adverse effect on reproduction or on the sexual satisfaction of the woman or her partner’.¹

Arora and Jacobs argue that such procedures should be considered permissible in young girls, so long as they are requested by the parents. Problematically, however, the only available studies assessing

reproductive and sexual outcomes associated with this particular intervention stem from surgeries performed on consenting adult women (or older adolescents); and even then the data are woefully incomplete.⁴⁹ But—again—let us just assume that, one day, researchers do produce a robust and convincing benefit-to-risk (or benefit-to-harm)⁵⁰ profile for non-therapeutic labiaplasty or similar procedures carried out on adults. Even so, we could not be sure that the profile would apply to children.³² For one thing, sexual outcomes, in particular, are highly subjective;⁵¹ and the women who undergo procedures akin to labiaplasty and subsequently report on their effects are unlikely to be representative of the general female population. This is because, overwhelmingly, they will have pursued their elective genital surgeries precisely because they were unsatisfied with their labia/vulva in some way, and wanted a change. In the case of a young girl, however, we cannot know, in advance, how she will (later) regard her own vulva—she might value her genitals in their intact state very highly—and so we are not in a position to judge how their modification would affect her sexual experience.

Sexual problems

Against this view, it might be argued that if genital tissue is excised early enough, a girl will not ‘know what she is missing’ when she reaches an older age. Similar arguments are raised in support of performing male circumcision as early as possible. But this does not guarantee a ‘harmless’ outcome. On the contrary, such an intervention could just as easily lead to feelings of loss or resentment—whether or not the surgery itself was consciously remembered. After all, a woman might reasonably wonder what sex, masturbation, and so on would have been like had her vulva been left intact, and feel angry that she was not given the chance to find out. In line with this perspective, there are indeed many documented reports of women (as well as men and intersex people) who had their genitals modified for non-therapeutic reasons in early childhood, who do in fact experience anger and resentment. These feelings, in turn, have the potential to impact negatively on sexual experience, quite apart from any ‘purely’ physical effects that would ensue from the loss of sensitive tissue (see [Box 1](#)).^{22 27 52–56}

The lesson here is that classification of childhood genital surgeries based on ‘predicted’ effects on sexuality is a mistake.

Box 1 What are the likely effects of procedures resembling labiaplasty on a woman’s sexual experience?

As Runacres and Wood note, ‘the labia minora are highly innervated along the entire free edge, and are involved in the process of engorgement during sexual arousal. It follows therefore that labiaplasty has the potential to remove tissue that contributes to sensory sexual arousal’. In fact, ‘the labia minora are second only to the clitoris for both sensation and sensitivity and are more sensitive than the vaginal introitus’.⁴⁹ In addition, the labia can be orally and manually manipulated, which may yield particular sensations that would be physiologically impossible if the tissue were removed—an outcome that Arora and Jacobs seem to regard as irrelevant. Whether such an outcome is on balance negative, of course, cannot be ‘scientifically’ determined; rather, it depends on an individual’s sexual preferences. For example, for those for whom the ability to fondle, etc., the labia is an important part of their sexual activity, the surgical reduction or elimination of this tissue would indeed be expected to ‘have an adverse effect on ... sexual satisfaction’. The upshot is that everyone is different. Thus, as Johnsdotter has argued, there is no consistent relationship between type or degree of genital cutting—whether in females, males or intersex people—and subjective sexual pleasure later on. Hence: ‘the current academic focus on the role of genitalia in understanding sexual pleasure is a dead end. While genitalia usually are central to sexual activity, and can be seen as a prerequisite for sexual intercourse, it is a misapprehension to see the state of them (cut or uncut) as determinative of the individual’s experience of the sexual encounter. Consequently, ongoing debates on enhancing or detrimental effects of circumcision practices often miss the point, since they are based in too narrow a theoretical framework of sexual pleasure’.⁵⁹ [p. 262]. See the online supplementary appendix for further discussion: <http://dx.doi.org/10.1136/medethics-2015-103030>.

Since everyone’s genitals are unique (including the specific distribution of nerve endings, how sensitive the tissue is in different parts, and so on), and since

people have different attitudes towards intact versus modified genitals, as well as different sexual preferences that can range rather widely, it is not ethically useful to make bland, medicalised statements about expected sexual satisfaction considered as an ‘average’ effect of some intervention. This is especially the case given that such an outcome is typically measured by reductive ‘scientific’ surveys involving numbered scales or similar, often failing to control for socially desirable responding.^{32 57}

Indeed, Arora and Jacobs’s willingness to rely on such statements in their endorsement of both male³¹ and female¹ forms of non-therapeutic genital alteration, rather than proceeding cautiously and recognising the limitations—or even nonexistence—of the pertinent data,³² is worrying. When a child’s genitals are on the line, along with the prospect of diminishing his or her future sexual function and/or satisfaction (the latter of which is a highly subjective notion), we should be extremely careful about making assumptions of no harm when the available data are so weak and contentious.

Cultural problems

A further problem is that Arora and Jacobs’s proposal is culturally incoherent. On the one hand, they decry ‘culturally insensitive’ laws and language that seek to eliminate female genital ‘mutilation’. But on the other hand, they offer a made-up, ‘medicalised’ ritual of their own design, without explaining how this would satisfy the real-life cultural impulses of the families who endorse FGA. As they themselves state, ‘We are ... not suggesting that people whose beliefs or sense of propriety leads them to perform these procedures on their children would necessarily accept alterations to their practices to conform to the authors’ views of what is acceptable’.¹ Indeed—why would they? As Hirsir Ali has written: ‘To understand this problem, we need to begin with parental motives. The ‘nicking’ option is regarded as a necessary cleansing ritual ... the clitoris is considered to be an impure part of the girl-child and bleeding it is believed to make her pure and free of evil spirits’. Problematically, however, for girls in at least some communities, the point of FGA is ‘to ensure their virginity ... and to curb their libido to guarantee sexual fidelity after marriage’. Thus, when FGA is intended ‘to ensure chastity before marriage and to curb female libido ... the nick option [would not be] sufficient’ from a cultural perspective.³⁷

Political problems

Finally, there is almost no chance that Arora and Jacobs’s suggestion will actually be taken up by lawmakers: it is a political non-starter. This can be inferred from the debacle that ensued in 2010, when the AAP issued a policy similar to the one proposed by the authors, in the name of ‘cultural sensitivity’. As Arora and Jacobs discuss, it was summarily retracted (just 1 month later) after it was met with fierce opposition and even outrage.⁵⁸ Opponents of the AAP policy included survivors of FGA such as Ayaan Hirsi Ali³⁷ and Soraya Mire;^{58–60} members of its own ranks;^{44 61 62} distinguished doctors from other countries;⁶³ US lawmakers;⁶⁴ and organisations such as Equality Now, an international advocacy network fighting to end female genital cutting.⁶⁵ In fact, I expect that Arora and Jacobs’s own proposal (as well as perhaps the *Journal of Medical Ethics*, for publishing it) will be met with a similar outcry. While their article constitutes an interesting (if unoriginal)³ academic exercise, Western societies have, for the most part, reached the seemingly irreversible conclusion that sharp objects should not be taken to the vulvas of little girls, unless it is to save their life or health.¹¹ Importantly, many immigrant communities in these same societies—derived from populations where FGA has traditionally been performed—have reached the same conclusion, and have willingly given up the practice.²⁶ It would be sobering for those who have struggled towards this end to see such progress reversed.

There are numerous other problems with Arora and Jacobs’s proposal, including misleading (or inaccurate) citations concerning key empirical assertions, but I do not have the space to cover them all in detail in this response. Instead, I refer the reader once again to the online supplementary appendix accompanying this

³⁷Similar patterns have played out in other countries. For example, as Leye *et al*³⁸ [p. 367] report (internal references omitted): ‘In 1992, two researchers in the Netherlands submitted a report to the Dutch Ministry of Health in which they proposed that as a step toward the total eradication of the practice, incisions of the clitoris be allowed in cases where the parents or family wanted to circumcise a girl. This kind of incision was considered as a nonmutilating form of ‘female circumcision’, as FGM is commonly referred to in the Netherlands. This report’s recommendation provoked a public debate about the issue that resulted in the total rejection of any form of FGM and, moreover, the rejection of any attempts to differentiate between mutilating and nonmutilating forms of ‘female circumcision’.

article, in which I discuss these and other matters (see <http://dx.doi.org/10.1136/medethics-2015-103030>).

CONCLUSION

In summary, if Western societies were to change their laws in order to allow for more ‘minor’ forms of FGA, this would likely result in: (a) disturbances and inconsistencies throughout their legal systems, possibly requiring new definitions of bodily assault and opening the door for inadvertent legal protection of a wide range of potentially harmful practices (typically carried out on children, who cannot adequately defend themselves); (b) removal of an important tool that reformers from within the affected communities rely on to solve the ‘collective action’ problem introduced by FGA; (c) regulatory challenges in tracking and monitoring FGA cutting sessions to ensure that they were not being used as opportunities for more invasive procedures; (d) exposure of young girls to an unknown amount of surgical risk in the absence of medical need, thereby placing doctors in an untenable position with respect to their professional duties; (e) accusations of cultural insensitivity, on account of outsiders having invented a ‘compromise’ intervention to be carried out in a medicalised setting, with the aim of replacing the traditional rituals that carry meaning for the relevant communities; and (f) widespread outrage among women who consider themselves victims and/or survivors of FGA as well as their allies, and other forms of political backlash.

In light of these considerations, let us return to Arora and Jacobs’s major conditional argument, namely, that ‘a liberal society that tolerates expression of culture and/or religion in the manner of male circumcision should also permit certain *de minimis* [female genital altering] procedures’.¹ I have tried to show that such permissiveness would result in a fiasco, making this suggestion (for all intents and purposes) a *reductio ad absurdum*. Accordingly, there is a growing trend among scholars of genital cutting, particularly in the fields of bioethics and law, of arguing that it is time to consider a less lenient position towards MGA (whether by banning it with a possible exception for sincere religious belief, or by regulating it in other ways), rather than a more lenient position towards FGA.^{8 9 17 43 66–68} As Arora and Jacobs themselves point out,

‘We acknowledge that issues of cultural sensitivity and gender discrimination in

the disparate treatment of male circumcision and FGA could also be treated by proscribing both, instead of the position for which we are advocating. In fact, many have criticised male circumcision as a human rights violation due to the lack of autonomous decision-making and the irreversible nature of the procedure'.¹

Whether it is in fact a fundamental human rights violation, or rather a morally objectionable practice on other grounds, 'it is clear that the current *laissez faire* attitude towards MGA that is typical of Western societies—and in particular the United States—can no longer be maintained without facing serious scrutiny. FGA and MGA are both highly problematic practices, with far more overlap between them (both physically and symbolically) than is commonly understood: they should not be discussed, therefore, in hermetically sealed moral discourses'.⁴ My own perspective is that the most promising way forward would be to argue for an 'autonomy-based' ethical framework,^{4, 9} whereby 'Children of whatever [sex or] gender should not have healthy parts of their most intimate sexual organs removed, before such a time as they can understand what is at stake in such a surgery and agree to it themselves'.⁶⁹

Acknowledgements Thanks are due to Mikey Dunn for constructive feedback on an earlier draft of this response.

Competing interests None declared.

Provenance and peer review Commissioned; internally peer reviewed.

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/medethics-2015-103030>).



CrossMark

To cite Earp BD. *J Med Ethics* 2016;**42**:158–163.

Received 27 November 2015

Accepted 2 December 2015

Published Online First 20 January 2016



- <http://dx.doi.org/10.1136/medethics-2014-102375>
- <http://dx.doi.org/10.1136/medethics-2015-103027>
- <http://dx.doi.org/10.1136/medethics-2015-103029>
- <http://dx.doi.org/10.1136/medethics-2016-103376>

J Med Ethics 2016;**42**:158–163.

doi:10.1136/medethics-2015-103030

REFERENCES

1 Arora K, Jacobs A. Female genital alteration: a compromise solution. *J Med Ethics* 2016;**42**:148–54.

2 WHO. Classification of female genital mutilation. (cited 12 Nov 2015). <http://www.who.int/reproductivehealth/topics/fgm/overview/en/>

3 Davis DS. Male and female genital alteration: a collision course with the law? *Health Matrix* 2001;**11**:487–570.

4 Earp BD. Female genital mutilation and male circumcision: toward an autonomy-based ethical framework. *Medical Bioethics* 2015;**5**:89–104.

5 Earp BD. Between moral relativism and moral hypocrisy: reframing the debate on "FGM." *Kennedy Inst Ethics J* 2016 (in press). https://www.academia.edu/10197867/Between_moral_relativism_and_moral_hypocrisy_Reframing_the_debate_on_FGM_

6 Svoboda JS, Darby R. A rose by any other name? Symmetry and asymmetry in male and female genital cutting. In: Zabus C, ed. *Fearful symmetries: essays and testimonies around excision and circumcision*. Amsterdam and New York: Rodopi, 2008:251–302.

7 Caldwell JC, Orubuloye IO, Caldwell P. Male and female circumcision in Africa from a regional to a specific Nigerian examination. *Soc Sci Med* 1997;**44**:1181–93.

8 DeLaet DL. Framing male circumcision as a human rights issue? Contributions to the debate over the universality of human rights. *J Hum Rights* 2009;**8**:405–26.

9 Svoboda JS. Promoting genital autonomy by exploring commonalities between male, female, intersex, and cosmetic female genital cutting. *Global Discourse* 2013;**3**:237–55.

10 Earp BD. Do the benefits of male circumcision outweigh the risks? A critique of the proposed CDC guidelines. *Front Pediatr* 2015;**3**:1–6.

11 Earp BD. The ethics of infant male circumcision. *J Med Ethics* 2013;**39**:418–20.

12 Earp BD. Things I've learned (so far) about how to do practical ethics. *Practical Ethics* 2014. <http://blog.practicaethics.ox.ac.uk/2014/03/things-ive-learned-so-far-about-how-to-do-practical-ethics/>

13 Waldeck S. Using circumcision to understand social norms as multipliers. *Univ Cincinnati Law Rev* 2003;**72**:455–526. <http://www.cirp.org/library/legal/USA/waldeck1/>

14 Ben-Yami H. Circumcision: what should be done? *J Med Ethics* 2013;**39**:459–62.

15 Krása K. Human rights for women: the ethical and legal discussion about female genital mutilation in Germany in comparison with other Western European countries. *Med Health Care Philos* 2010;**13**:269–78.

16 Rahman A, Toubia N. *Female genital mutilation: a practical guide to worldwide laws and policies*. London: Zed Books, 2000.

17 Merkel R, Putzke H. After Cologne: male circumcision and the law. Parental right, religious liberty or criminal assault? *J Med Ethics* 2013;**39**:444–9.

18 Boyle GJ, Svoboda JS, Price CP, et al. Circumcision of healthy boys: criminal assault? *J Law Med* 2000;**7**:301–10. http://epublications.bond.edu.au/cgi/viewcontent.cgi?params=/context/hss_pubs/article/1121/type/native&path_info=

19 Chegwidden J. The assault we ignore. Children's Rights in Europe: Recent Developments. 17 Sept 2013. <https://vimeo.com/78999394>

20 Svoboda JS, Van Howe RS, Dwyer JG. Informed consent for neonatal male circumcision: an ethical and legal conundrum. *J Contemp Health Law Policy* 2000;**17**:61–133.

21 Adler P. Is circumcision legal? *Richmond J Law Public Interest* 2013;**XVI**(iii):439–83.

22 Ford K. "First, Do No Harm": the fiction of legal parental consent to genital-normalizing surgery on intersexed infants. *Yale Law Policy Rev* 2001;**19**:469–88.

23 Svoboda JS, Adler P, Van Howe RS. Circumcision is unethical and unlawful. *J Law Med Ethics* (in press).

24 Blackstone W. *The commentaries of Sir William Blackstone, Knight, on the Laws and Constitution of England*. 1796 edn. Chicago, Illinois: ABA Publishing, 2009.

25 Ludbrook R. The child's right to bodily integrity. *Curr Issues Crim Justice* 1995;**7**:123–32. http://heinonline.org/HOL/Page?handle=hein.journals/cicj7&div=23&g_sent=1&collection=journals

26 Johnsdotter S, Essén B. Cultural change after migration: circumcision of girls in Western migrant communities. *Best Pract Res Clin Obstet Gynaecol* 2015 (in press):<http://www.sciencedirect.com/science/article/pii/S1521693415001959>

27 Norman K, Hemmings J, Hussein E, et al. *FGM is always with us: experiences, perceptions, and beliefs of women affected by female genital mutilation in London*. Options Consultancy Services and FORWARD, 2009. <http://www.forwarduk.org.uk/563/>

28 Mackie G. Ending footbinding and infibulation: a convention account. *Am Sociol Rev* 1996;**61**:999–1017.

29 Mackie G. Female genital cutting: the beginning of the end. In: *Female "Circumcision" in Africa: Culture, Controversy and Change*. Boulder, CO: Lynne Reinner Publishers, 2000:253–82.

30 Everett JA, Earp BD. A tragedy of the (academic) commons: interpreting the replication crisis in psychology as a social dilemma for early-career researchers. *Front Psychol* 2015;**6**:1–4.

31 Jacobs AJ, Arora KS. Ritual male infant circumcision and human rights. *Am J Bioeth* 2015;**15**:30–9.

32 Earp BD. Sex and circumcision. *Am J Bioeth* 2015;**15**:43–5.

33 Ungar-Sargon E. On the impermissibility of infant male circumcision: a response to Mazor (2013). *J Med Ethics* 2015;**41**:186–90.

34 Darby RJ. The child's right to an open future: is the principle applicable to non-therapeutic circumcision? *J Med Ethics* 2013;**39**:463–8.

35 Carmack A, Notini L, Earp BD. Should surgery for hypospadias be performed before an age of consent? *J Sex Res* 2015 (in press). <http://www.tandfonline.com/doi/full/10.1080/00224499.2015.1066745>

36 Maslen H, Earp BD, Cohen Kadosh R, et al. Brain stimulation for treatment and enhancement in children: an ethical analysis. *Front Hum Neurosci* 2014;**8**:1–5.

37 Hirsir Ali A. Why are American doctors mutilating girls? *The Daily Beast*, 2010. <http://www.thedailybeast.com/articles/2010/05/20/ayaan-hirsir-ali-on-injustice-of-female-genital-mutilation.html>

38 Leye E, Powell RA, Nienhuis G, et al. Health care in Europe for women with genital mutilation. *Health Care Women Int* 2006;**27**:362–78.

39 Williams N, Kapila L. Complications of circumcision. *Br J Surg* 1993;**80**:1231–6.

40 Krill AJ, Palmer LS, Palmer JS. Complications of circumcision. *Scientific World J* 2011;**11**:2458–68. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3253617/>

41 Baskin L, Canning D, Snyder H, et al. Treating complications of circumcision. *Pediatr Emerg Care* 1996;**12**:62–8.

42 American Academy of Pediatrics Task Force on Circumcision. Male circumcision. *Pediatrics* 2012;**130**:e756–85.

43 Dustin M. Female genital mutilation/cutting in the UK: challenging the inconsistencies. *Eur J Womens Stud* 2010;**17**:7–23.

44 Van Howe RS. The American Academy of Pediatrics and female genital cutting: when national organizations are guided by personal agendas. *Ethics Med* 2011;**27**:165–74.

45 Catania L, Abdulcadir O, Puppo V, et al. Pleasure and orgasm in women with Female Genital Mutilation/Cutting (FGM/C). *J Sex Med* 2007;**4**:1666–78.

46 Donovan B. Review of "In Favour of Circumcision" by Brian Morris. *Venerology* 1999;**12**:68–9.

- 47 Earp BD, Darby R. Does science support infant circumcision? *The Skeptic* 2015;25:23–30. https://www.researchgate.net/publication/269899744_Does_science_support_infant_circumcision
- 48 Goldman R. Cultural bias may contribute to flawed policy. *Pediatrics* 2010;125. <http://pediatrics.aappublications.org/content/125/5/1088.comments#cultural-bias-may-contribute-to-flawed-policy>
- 49 Runacres SA, Wood PL. Cosmetic labiaplasty in an adolescent population. *J Pediatr Adolesc Gynecol* 2015 (in press). <http://www.ncbi.nlm.nih.gov/pubmed/26453828>
- 50 Darby R. Risks, benefits, complications and harms: neglected factors in the current debate on non-therapeutic circumcision. *Kennedy Inst Ethics J* 2015;25:1–34.
- 51 Johnsdotter S. Discourses on sexual pleasure after genital modifications: the fallacy of genital determinism (a response to J. Steven Svoboda). *Global Discourse* 2013;3:256–65.
- 52 Darby R, Cox L. Objections of a sentimental character: the subjective dimensions of foreskin loss. In: Zabus C, ed. *Fearful symmetries: essays and testimonies around excision and circumcision*. Amsterdam and New York: Rodopi, 2009: 0145–68.
- 53 Watson L. *Unspeakable mutilations: circumcised men speak out*. New Zealand: Amazon Media, 2014.
- 54 Bossio J. Examining sexual correlates of neonatal male circumcision. Doctoral dissertation, 2015. <https://qspace.library.queensu.ca/handle/1974/13627>
- 55 Foldès P, Cuzin B, Andro A. Reconstructive surgery after female genital mutilation: a prospective cohort study. *Lancet* 2012;380:134–41.
- 56 Dreger A. *Galileo's middle finger: heretics, activists, and the search for justice in science*. New York: Penguin Press, 2015.
- 57 Earp BD. The need to control for socially desirable responding in studies on the sexual effects of male circumcision. *PLoS ONE* 2015;10:1–12. https://www.researchgate.net/publication/281976691_The_need_to_control_for_socially_desirable_responding_in_studies_on_the_sexual_effects_of_male_circumcision?ev=pub_cit_inc
- 58 Chen S. Pediatricians now reject all female genital cutting. 2010 [cited 24 Nov 2015]. <http://edition.cnn.com/2010/HEALTH/05/27/AAP.retracts.female.genital.cutting/>
- 59 Pollack M. Circumcision: gender and power. In: Denniston G, Hodges F, Milos M, eds. *Genital cutting: protecting children from medical, cultural, and religious infringements*. Dordrecht: Springer Netherlands, 2013:297–305.
- 60 Mire S. *The girl with three legs: a memoir*. Chicago: Chicago Review Press, 2011.
- 61 Strandjord S. Opportunity for education plus protection of minors. *Pediatrics* 2010;125. <http://pediatrics.aappublications.org/content/125/5/1088.comments#opportunity-for-education-plus-protection-of-minors>
- 62 Bamzai A. Re: AAP response to eLetters. *Pediatrics* 2010;125: <http://pediatrics.aappublications.org/content/125/5/1088.comments#re-aap-response-to-eletters>
- 63 Bewley S. Sincerity, retraction and apology? *Pediatrics* 2010;125. http://pediatrics.aappublications.org/content/125/5/1088.short/reply#pediatrics_el_50347
- 64 Luscombe B. Female genital cutting: will US doctors approve nicks? *Time* 2010. <http://content.time.com/time/health/article/0,8599,1988434,00.html>
- 65 Equality Now. American Academy Of Pediatrics (AAP) is advocating for US pediatricians to perform certain types of Female Genital Mutilation (FGM). Equality Now. 2010 (cited 24 Nov 2015). <http://www.equalitynow.org/american-academy-pediatrics-aap-advocating-us-pediatricians-perform-certain-types-female-genital-mut>
- 66 Fox M, Thomson M. Short changed? The law and ethics of male circumcision. *Int J Child Rights* 2005;13:161–81.
- 67 Tasmania Law Reform Institute. *Non-therapeutic male circumcision*, 2012. http://www.utas.edu.au/__data/assets/pdf_file/0006/302829/Non-Therapeutic-Circ_Final-Report-August-2012.pdf
- 68 Hellsten SK. Rationalising circumcision: from tradition to fashion, from public health to individual freedom—critical notes on cultural persistence of the practice of genital mutilation. *J Med Ethics* 2004;30:248–53.
- 69 Earp BD. Female Genital Mutilation (FGM) and male circumcision: should there be a separate ethical discourse? *Practical Ethics* 2014 (cited 24 Nov 2015). <http://blog.practicaethics.ox.ac.uk/2014/02/female-genital-mutilation-and-male-circumcision-time-to-confront-the-double-standard/>

Appendix to: “In defence of genital autonomy for children”

Brian D. Earp

Visiting Scholar, The Hastings Center Bioethics Research Institute,
Garrison, NY, USA

Introduction

This is an Appendix to the article, “In Defence of Genital Autonomy for Children”(1) which is itself a response to “Female Genital Alteration—A Compromise Solution” by Arora and Jacobs.(2)

The Appendix is divided into two parts. In the first part (Part I), I present an overview of various “overlaps,” both physical and symbolic, between male and female forms of non-therapeutic genital alteration (MGA, FGA). This is to show why the conditional argument by Arora and Jacobs—namely that *if* it is the case that MGA should be widely tolerated, then so should some forms of FGA—is valid even if it is not, as I argue in my response piece, sound. In the second part (Part II), I provide a point-by-point response to various specific claims by Arora and Jacobs that are misleading, inaccurate, or otherwise problematic, but which I did not have the space to cover in the main article.

PART I. A brief overview of the morally-relevant “overlaps” between FGA and MGA

In this section, I provide a brief sketch of the “overlaps” between non-therapeutic male and female genital altering procedures. I recognize that many of the claims in this section will seem counterintuitive to those who are new to this area given the prevailing popular and academic discourses; I can only hope that the reader will take the time to follow up on the references I cite in support of them. For a much more sustained defense of the view presented in this section, please see my essay, “Female Genital Mutilation and Male Circumcision: Toward an Autonomy-Based Ethical Framework.”(3)

Area of overlap #1: harm

Both male and female forms of non-therapeutic genital alteration fall on a wide spectrum. The physical and sexual harms that they may cause substantially overlap, depending upon the type of procedure, how much tissue (and of what kind) is removed, which instruments are used and how

sterile they are, what level of training the practitioner has, and so on. In particular, several forms of WHO-recognized FGM procedures, such as cutting and/or removing part(s) of the clitoral hood (such as is common in Malaysia), as well as some interventions into the labia, are uncontroversially less physically harmful than the most widespread forms of male circumcision. Conversely, several forms of male genital cutting/alteration, such as subincision in parts of aboriginal Australia, or tribal circumcision as it is carried out among the Xhosa of South Africa, can be at least as physically harmful as the most widespread forms of FGM, and are frequently much more harmful as a matter of practice.(3)

Area of overlap #2: benefits

The benefits of both male and female forms of non-therapeutic genital alteration overlap as well. As I have argued elsewhere,(3,4) the health benefits that have been attributed to male circumcision, at least in a Western context, are relatively minor (in terms of absolute risk reduction for various diseases, many of which can be prevented and/or treated non-surgically); they apply mostly after an age of sexual debut (i.e., when meaningful consent could at least in principle be obtained); they can be achieved in much less harmful ways (such as by the practice of basic hygiene and the adoption of safe sex practices); and, according to the balance of medical opinion on a global scale, they do not very likely outweigh the risk of immediate complications plus delayed complications and longer-term harms(5) (contrary to the insinuations of Arora and Jacobs; see the discussion in Part II of this Appendix).

On the other side of the coin, we do not actually know that certain minor forms of FGA—such as neonatal labiaplasty, or perhaps other interventions—would not confer the same degree of prophylaxis against disease as male circumcision purportedly does, in virtue of removing healthy tissue from the vulva that could later host an infection, cancer, or other malady. Indeed, it is illegal, in Western countries, to conduct a trial to find this out.(4) That said, there is in fact some observational evidence that FGA may be associated with certain health benefits, including reduced transmission of HIV.(3)

With respect to purported socio-cultural and/or religious (or spiritual) benefits, both male and female forms of non-therapeutic genital alteration are defended in largely similar terms by those who endorse them.(3,6–13) Speaking generally, as Gunning notes, “Both can be seen as unnecessary alterations of normal, healthy genitalia justified by questionable health benefits and

bolstered by culturally, socially, or religiously defined notions of aesthetics and clearly delineated binary ideas of gender.”(14)[p. 655-656]

Area of overlap #3: symbolic meanings

Contrary to common wisdom, FGA “is not always associated with, nor a reflection of, sexist and patriarchal norms; nor are the norms associated with male genital cutting always as morally innocent as is typically assumed.”(3) For example, among the Kono of Sierra Leone, “there is no cultural obsession with feminine chastity, virginity, or women’s sexual fidelity ... because the role of the biological father is considered marginal and peripheral to the central ‘matricentric unit.’”(15) In this context (as in many other African contexts) “male and female genital alterations are performed in parallel ceremonies, are not primarily intended to reduce sexual pleasure, and the operations are seen as mirror images of each other.”(3)

At the same time, as Fox and Thomson show, certain other forms of male genital cutting (as well as female genital cutting) can in fact be understood as gendering practices tied to patriarchal notions of masculinity, as well as to cultural systems based on male privilege and the ritual exclusion of women.(16) For example, as Kimmel argues, male circumcision “means [the] reproduction of patriarchy.” In the Jewish tradition, “Abraham cements his relationship to God by a symbolic genital mutilation of his son. It is on the body of his son that Abraham writes his own beliefs. In a religion marked by the ritual exclusion of women, such a marking not only enables Isaac to be included within the community of men [but] he can also lay claim to all the privileges to which being a Jewish male now entitles him.” To circumcise one’s son, therefore, is “to accept as legitimate 4000 years [of] patriarchal domination of women.”(17)

Finally, as Arora and Jacobs themselves point out (and as many others have argued), the supposed distinction between “religion” and “culture” as a means of distinguishing male vs. female forms of ritual genital cutting is logically untenable, historically inaccurate, and anthropologically ill-informed.(6,18,19)

Implication

Taken together, these overlaps imply that “a stark moral distinction between male and female forms of non-therapeutic, non-consensual genital alteration may be impossible to maintain on principled grounds.”(3) As a consequence of this fact, liberal societies face a choice if they wish

to be consistent.(20) Roughly speaking, they can either allow both types of genital alteration to be carried out on similar terms, or they can disallow (or at least discourage) both types of genital alteration on similar terms.

Arora and Jacobs prefer the former option; I prefer the latter. In the next section, I take a closer look at the arguments and evidence presented by Arora and Jacobs in support of their position, which I argue is ultimately untenable. Here I go beyond what I was able to cover in my original response piece.

Part II. A closer look at the various claims advanced by Arora and Jacobs

In this section I address specific weaknesses in Arora and Jacobs's (A&J's) argument, by giving a point-by-point response. I will move sequentially through their paper, and the format will be to give a quote from A&J, followed by a discussion of any problems contained therein.

“The World Health Organization (WHO), American Academy of Pediatrics (AAP), and the American Congress of Obstetricians and Gynecologists (ACOG) have policies in place to support circumcision.”

There are several problems with this statement. First, the WHO has a policy in place to support so-called Voluntary Medical Male Circumcision (VMMC), not in “Western” countries (which is the main subject of A&J's proposal), but rather in sub-Saharan Africa as a form of partial prophylaxis against heterosexually transmitted HIV (female-to-male only) in areas with high base-rates of such transmission, and low base rates of male circumcision.(21) Crucially, the “Voluntary” part of VMMC implies (or should imply) that the circumcision should be undertaken by an individual who can provide his own consent, not an infant or young child (who cannot). Nobody objects to truly informed, consensual adult circumcision; it is the circumcision of infants and children that is ethically contentious. Moreover, the data that the WHO appeals to in support of its policy were derived from Randomized Control Trials (RCTs) performed with adult male participants (in Africa), not infants or young children (in Western countries). Since “the spread of disease, including sexually transmitted infections, is determined much more by socio-behavioral and situational factors than by strictly anatomical-biological factors, such as the presence or absence of a foreskin ... the apparent findings from these studies cannot be simply mapped on to non-analogous public health environments, nor to circumcisions performed earlier in life, i.e., before an age of sexual debut.”(4)

Second, A&J do not mention that the policy by the AAP they cite (which does not recommend circumcision, but rather leaves the decision up to parents) was roundly criticized by international pediatric and sexual health experts,(22) and is in fact inconsistent with comparable policies from many other pediatric societies in peer nations, including the recently-released statement by the Canadian Paediatric Society.(23,24) As Wolfram Hartmann, President of the German Pediatric Society, explained, “The statement from the AAP ... has been graded by almost all other paediatric societies and associations worldwide as being scientifically untenable.”(25) For an incomplete sample of published critiques of the 2012 AAP policy statement and technical report—none of which were cited by Arora and Jacobs in their paper—see the references collected here.(4) For an incomplete sample of medical bodies which, in their current policies, either fail to conclude that the benefits of involuntary male circumcision outweigh the risks, or state that the risks in fact outweigh the benefits, see statements from: the Canadian Paediatric Society;(23) the Royal Australasian College of Physicians;(26) the Royal Dutch Medical Association;(27) the German Academy for Pediatrics and Adolescent Medicine;(25) the Swedish, Norwegian, and Icelandic Pediatrics Societies;(28) and the National Health Service of England.(29) Why Arora and Jacobs decided to emphasize the position of a minority of medical bodies—from the single developed country in the world that practices routine circumcision for non-religious reasons—whose policies could be interpreted as supporting their views on health benefits for male circumcision is unclear.

Finally, please note that the ACOG does not have its own policy “supporting” male circumcision, but rather signed on to the AAP policy as a formal gesture. The document available from the ACOG’s own website actually states: “At the present time, there is not enough information to recommend routine newborn circumcision for health reasons.”(30)

“While years of advocacy and legislation aimed at eliminating non-therapeutic procedures on female external genitalia has resulted in a decline in the prevalence of the practice, the magnitude of this decline has been soberingly small. ... In a study in Somalia, the country in the world with the highest prevalence of these procedures, 81% of subjects underwent infibulation and only 3% did not have FGA. Eighty-five percent had an intention to subject their daughters to an extensive FGA procedure, and 90% supported the continuation of the practice.

This citation is misleading. The study involved a sample of 215 persons from one county in one town in Somalia; there is no way of knowing whether the reported estimates are nationally

representative.(31) Moreover, the statement that “Eighty-five percent” of participants “had an intention to subject their daughters to an extensive FGA procedure” is false: as the authors of the report themselves stated, of this 85%, fully “154 (71.6%) were planning to use the *Sunna* form [i.e., the least extensive form], with the reason behind their decision either being because they considered the *Sunna* form to be harmless or because it was seen as a religious requirement.”(31)[p. 4]. Similarly, with respect to the statement (from A&J) that “90% supported the continuation of the practice,” it should be noted that the authors of the report actually wrote: “Of this, 164 (76.3%) persons supported the continuation of the Sunna form only, while [only] 10.7% supported the continuation of all forms” of the practice.(31)[p. 4] The broader point, however, is that there has, in fact, been significant progress in many African countries in reducing the rate of non-therapeutic FGA procedures (as A&J themselves concede later on in the paragraph).

Even more importantly, however, what A&J appear to be advocating (although their language is ambiguous on this point) is a change in laws in Western countries, not Africa. So the only data that are relevant to their argument is the prevalence of FGA in Western countries (among immigrant populations), and the effect of laws that prohibit FGA on its prevalence in those countries. In other words, it is a non-sequitur to argue that, since the effects of anti-FGA advocacy in African countries have been “soberingly small” (whether or not that really is the case), we should therefore “relax” the laws concerning FGA in countries like the United States!

On this point, A&J provide no compelling estimates of FGA prevalence in Western societies, but instead state that: “Immigrants to Western nations may continue to subject their daughters to genital alteration, though the frequency is difficult to assess.” By contrast, in a recent review of the evidence, Johnsdotter and Essén report that: “Exploratory studies show trends of radical change of this harmful practice, especially the most extensive form of its kind. The widespread interpretation that Islam would require circumcision of girls becomes questioned when, e.g., Somalis meet other migrants, such as Arab Muslims, who do not circumcise their daughters. The few criminal court cases for circumcision of girls that have taken place in Western countries corroborate the conclusion that substantial change of the practice has occurred among migrants.”(32) Insofar as A&J’s policy proposal really is motivated by a concern about “widespread” FGA being carried out on an “underground” basis in Western countries, due to their restrictive laws concerning the practice, it is not clear that there is enough of a problem that

changes to these laws should in fact be seriously entertained.

“While laws enacted in [Western] societies make procedures that alter a female’s external genitalia illegal, they may in some instances worsen health outcomes by driving the practice underground, by sending female children to Africa, or by inviting circumcisers to the West.”

They may have this effect, or they may not. Unfortunately, the authors do not provide any evidence that prohibitions on FGA in Western countries lead to worse health outcomes overall by, e.g., “driving the practice underground.” Instead, they cite a paper by Leye et al., which in turn cites another paper by Leye (and a colleague) in which those authors state: “The true magnitude of the problem of FGM in [the studied countries] could not be assessed, and this lack of accurate FGM prevalence data make it hard to substantiate the claim for services, legislation and funds for prevention work.”(33)[p. 47] It is easy to speculate. But on a harm-reduction analysis, prohibition would only be an imprudent strategy if the amount of harm that was added through the postulated “underground” FGA procedures (including overseas trips, etc.) as a consequence of the law were greater than the amount of harm that was reduced by an overall drop in prevalence of FGA (due to its being illegal).

In short, we do not know how many girls are at risk of FGA in Western countries, and we have no evidence that the law (as it stands) is not an effective harm-reduction measure compared to alternatives (including A&J’s proposal), particularly when it comes to “health outcomes.”(32) In Africa and the Middle East, by contrast, where many communities see the prohibitions in their countries as being effectively forced on them by cultural outsiders/ former colonial powers, the laws may indeed lead to resentment and backlash and may therefore be much less effective. But this would suggest that cultural imperialism is a problem,(34) not (necessarily) that the laws in Western countries should be changed to accommodate the practices of immigrants who come to their shores (just as we would not expect, e.g., China to change its laws to accommodate the controversial practices of Westerners who might immigrate there). As Chapin (quoted in Belluck)(35) stated in response to an earlier U.S. policy proposal (by the AAP) similar to the one advanced by A&J: “There are countries in the world that allow wife beating, slavery and child abuse, but we don’t allow people to practice those customs in this country. We don’t let people have slavery ‘a little bit’ because they’re going to do it anyway, or beat their wives ‘a little bit’ because they’re going to do it anyway.” More generally, when certain acts in a host country are considered morally impermissible, such as sexual assault—yet nevertheless persist for whatever

reason (whether in immigrant communities or in the majority population)—this does not normally cause us to consider “relaxing” the laws against sexual assault (to repeat that example), including its “minor” forms.

Now, one could certainly argue that the law should not be the only tool for reducing harm associated with FGA, especially if it is being applied in a “sensationalized, ethnocentrist, racist, culturally insensitive, and simplistic” manner as A&J worry. But rather than “relaxing” the law, the law could instead be complemented with, e.g., sensitivity training, and could be applied in a nonracist, non-ethnocentric way by being expanded to cover similar practices that happen to be more popular in the host country.(34)

“To accommodate cultural beliefs while protecting the physical health of girls, we propose a compromise solution in which liberal states would legally permit ‘de minimis’ FGA in recognition of its fulfillment of cultural and religious obligations, but would proscribe those forms of FGA that are dangerous or that produce significant sexual or reproductive dysfunction.”

First, there is a question of whether it is the role of liberal state to “accommodate cultural beliefs,” and a further question about what a “cultural belief” is in the first place, but those are minor points. More important is the suggestion that FGA should be tolerated if it is not “dangerous” or does not “produce significant sexual or reproductive dysfunction.” Language can be slippery. What do A&J mean by “significant” sexual dysfunction? Is there (by contrast) “insignificant” or “mild” sexual dysfunction that would still be permitted on this analysis? And who gets to decide what degree of sexual dysfunction is “significant” (and in what sense), or what degree of risk of such dysfunction is tolerable? One plausible argument is that it is the individual herself (whose possible sexual dysfunction is at stake) who should be able to decide the degree of risk she is willing to take on—as well as what she considers to be “significant” in this very personal area—when she reaches an age of mental competence. This is as opposed to the individual’s parents before such an age.

“Grouping all forms of FGA in discourse and condemnation assumes that all FGA procedures carry the same risks, which is medically inaccurate.”

This is not true. It is not in fact necessary to assume that all FGA procedures carry the same medical risks in order to object to them without discrimination, since “medical risk” is not necessarily the only morally relevant feature that they share. Indeed, most people who argue

against all forms of FGA acknowledge that there is more or less risk of certain kinds of physical or psychological harms depending upon the type of the procedure and the context in which it is carried out. What they suggest does not vary between type, however, and which may therefore justify their being “grouped together,” is that all forms are in conflict with (what they see as) (a) the child’s right to bodily integrity, and (b) her interest in making an informed decision about irreversible genital surgery when she understands what is at stake in such an intervention and is competent to weigh alternatives.(36)

“Authors arguing against all forms of FGA ... argue that physical well-being trumps social and cultural well-being.”

A&J do not cite any authors specifically, so it is hard to know whose work they are referring to. However, speaking generally, those who argue against all forms of FGA do not necessarily suggest that physical well-being trumps social and cultural well-being; rather, they include in “social and cultural well-being” a child’s interest in having her future autonomy preserved when it comes to making controversial and irreversible bodily alterations that she (or he, in the case of MGA) may later come to resent. Moreover, they question why a child’s social and cultural well-being should hinge on having a non-therapeutic genital surgery performed before an age of meaningful consent. As an alternative, they suggest that cultural groups should reconsider holding a child’s social well-being hostage to such a surgery, and should instead love and care for the child unconditionally until the child herself can make an informed decision about so “private” a part of her own body.(37–39)

“We believe that all procedures should be performed with adequate analgesia.”

What do A&J consider “adequate” in this regard? In the case of male circumcision, where pain control has been better studied, Bellieni et al. argue that “no procedure has been found to definitively eliminate pain; the gold standard procedure to make MC totally pain free has not yet been established.”(40) Just as with their vague phrasing about “significant sexual dysfunction,” then, one could reasonably wonder how much pain, exactly, A&J think is acceptable to inflict on a young girl for non-therapeutic reasons.

“Given that the more extensive forms of FGA are physically harmful and may constitute oppression towards women, these practices should be actively discouraged by means such as education, social pressure, regulation, and prohibition. [However, since] ‘de minimis’ alternate

procedures are not associated with the same risks of long-term harm, these should be encouraged as a compromise solution that upholds cultural and religious practices without sacrificing the health and well-being of female children.

A&J change their standard of analysis halfway through this paragraph. They begin by arguing that the “more extensive” forms of FGA should be prohibited because, in addition to being “physically harmful” they also “may constitute oppression towards women.” But then when it comes to the more “minor” forms that they believe should be “encouraged” (note that this is a stronger claim than that they should merely be “tolerated”), they suddenly forget the issue of oppression towards women. But these “minor” forms, too, may also constitute oppression towards women (and therefore be objectionable), even if they are not as “physically harmful” as the more extensive forms.(41)

“We use the term ‘procedure’ in the context of FGA rather than ‘surgery’ to emphasize that there is no medical benefit established by well-designed trials, and that the primary purpose is not health-related.”

A&J are indirectly alluding to male circumcision here, which is presumed to have medical benefits “established by well-designed trials,” but there are a couple of things wrong with this (implied) contrast. First, in the case of *ritual* male circumcision, which is the relevant comparison, the “primary purpose” is not “health-related” either. Health benefits are often referred to when secular critics question the practice, but that is also the case for FGA.(7,13,42) Second, there is no “medical benefit established by well-designed trials” (for FGA) because it is illegal to conduct such trials in the first place. There is in fact research showing a “significant and perplexing inverse association between reported female circumcision and HIV seropositivity,”(43) but as the medical anthropologist Kirsten Bell has noted (personal communication, January 16, 2015): “These findings, which were presented at an International AIDS Society conference in 2005, have never been published in a peer-reviewed journal and it is difficult to imagine any agency willing to entertain [the authors’] call for further research. Indeed, the topic is self-evidently a non-starter. Regardless of any evidence that might suggest an association, it is impossible to imagine a parallel research agenda [to the one on male circumcision] solidifying around the procedure, irrespective of whether the surgery was conducted in a medical context and [irrespective of] the extent of cutting involved.”

“Category 1 includes procedures that should almost never have a lasting effect on morphology or function if performed properly. A small nick to the vulvar skin fits into this category. Category 2 consists of procedures that create morphologic changes, but are not expected to have an adverse effect on reproduction or on the sexual satisfaction of the woman or her partner. Examples include surgical retraction of the clitoral hood or procedures resembling labiaplasty as performed in Western nations.”

Several points here. To count for Category 1 (on A&J’s proposed typology), the procedure must “almost never” have a lasting effect on morphology or function if performed properly. This raises the regulatory issue of how individuals will be trained (and by whom), and how “proper” performance will be monitored and assessed. In this context, we should ask: How many girls should we be willing to subject to “improperly performed” Category 1 procedures (which would in fact risk changing morphology and/or damaging function), as we sort out the issues of training and regulation? Here it should be remembered that even the slightest nick to the vulvar skin carries risk: the knife can slip; the equipment may not have been properly sterilized, introducing the risk of infection, and so on.

With respect to male circumcision, which has been performed in a Western context for centuries, and primarily by medical professionals, there is—even so—little guarantee of a “proper performance.” A recent study in the *Canadian Urological Association Journal*, for example, found: “Most physicians performing neonatal circumcisions in our community have received informal and unstructured training, [leading to] unsatisfactory results [being] witnessed in our pediatric urology practice. Many practitioners are not aware of the contraindications to neonatal circumcision and most non-surgeons perform the procedure without being able to handle common post-surgical complications.”(44) How much worse might the situation be with FGA procedures performed on a diminutive genital organ, given that comparable MGA procedures have had a much longer time period to become stabilized and thus “properly” performed?

For Category 2, A&J give the examples of surgical retraction of the clitoral hood and procedures akin to labiaplasty, arguing that these would “create morphological changes, but are not expected to have an adverse effect on reproduction or the sexual satisfaction of the woman or her partner.” It must first be noted that it is likely to be difficult to retract “just” the clitoral hood, since, in a young girl, the external clitoral organ (including its prepuce) is typically very small. Thus there would be a substantial risk of damaging (or over-exposing) the glans clitoridis, which is extremely

sensitive and which can be excruciating to touch directly (i.e., without the covering of the clitoral prepuce, which typically only retracts in a state of high arousal).(45) Why this is not seen as posing a meaningful risk of an “adverse effect on ... the sexual satisfaction of the woman” is unclear.

With respect to labiaplasty, it might be helpful to keep in mind what sort of tissue is at stake in this procedure, so that we can assess the likely effects of its removal on sexual sensation, function, etc.(46) As Schober et al. state: “Labia minora is highly innervated along its entire edge. Related vascular compartment tissue involved in engorgement during sexual arousal makes this tissue important for sexual response. Labiaplasty risks removal of tissue with an important contribution to sensory sexual arousal.”(47) Brzyski and Knudtson state: “The labia majora [are] relatively large, fleshy folds of tissue that enclose and protect the other external genital organs. They are comparable to the scrotum in males. The labia majora contain sweat and sebaceous glands, which produce lubricating secretions [while] the labia minora are lined with a mucous membrane, whose surface is kept moist by fluid secreted by specialized cells.”(48) Of course, if an adult female would like to sacrifice any of this tissue—with its concomitant sensory and protective functions, lubricating secretions, and so on—for “cosmetic” (or other) reasons, one could argue that that is her business. But to suggest that removing this tissue from a minor, who might very well grow up to wish that she still had her labia intact, is only a “*de minimis*” intervention, strikes me as somewhat harder to defend than the authors seem to think it is.

Indeed, it is worrying that A&J would include labiaplasty in a category that is “not expected to have an adverse effect on [the] sexual satisfaction of the woman or her partner” while providing no evidence whatsoever in support of this assertion. What is relevant to this debate is the likely (future) sexual effects of labiaplasty (or similar) as performed on a young girl. But since performing labiaplasty on young girls is illegal in Western societies, it is impossible to collect the relevant data. So, there is no way to know whether such an intervention would in fact fall into Category 2 unless it were first permitted, but A&J argue that it should be permitted on account of falling into Category 2. That seems somewhat circular.

But at least A&J are consistent. In a previous paper,(49)[p. 32] they argued that ritual infant male circumcision should have “little negative effect on sexual health and functioning,” based on data collected from trials of medicalized adult circumcision that involved only limited follow-up and which did not use validated questionnaires.(24) In a reply piece,(50) I pointed out that these data,

quite apart from being of poor quality, were not actually applicable (along several dimensions) to the form of circumcision A&J were defending. Moreover, I stressed that the foreskin itself consists of a substantial amount of touch-sensitive tissue that is both manually and orally manipulable, and that its removal necessarily eliminates any concomitant motions and sensations. To this, they replied: “Sexually sensitive skin unquestionably is removed during circumcision. It does not follow that this causes a loss of function or of satisfaction, or that remaining skin cannot compensate.”(51)[p. W1]

But it depends on how you define function, how you measure satisfaction—and who you ask. Any sexual functions (including masturbatory gestures) that require manipulation of the foreskin are, in fact, lost to circumcision. Similarly, any sexual functions involving manipulation of the labia (including some possible aspects of oral sex) are, in fact, lost to labial excision. Whether the inability to have one’s foreskin or labia manually or orally manipulated counts against one’s sexual satisfaction will depend upon one’s sexual preferences. Some men who have sex with men (MSM) use their foreskins in a sexual act known as “docking”(52) – an act that is impossible if the foreskin has been removed. In fact, in a recent study, “men indicated a strong preference toward intact penises for all sexual activities assessed and held more positive beliefs about intact penises.”(53) Similarly, the labia can be tugged, stretched, sucked on, and otherwise fondled during sexual interaction, and for those for whom such activities are an important part of their sexual experience, the loss of labia would indeed be a problem.

“Surgical resection of the clitoral hood is the vulvar procedure that most closely resembles male circumcision.”

The two interventions are similar in that they both involve resection of the genital prepuce. However, the highest concentration of sensitive nerve endings for the female is in the glans clitoris, not the clitoral prepuce; whereas, according to some researchers, in males the inverse is true: the glans penis is relatively lacking in fine-touch neuro-receptors, whereas the penile prepuce has a greater concentration of such receptors.(45) Moreover, the clitoral prepuce is very small, whereas the penile prepuce is very large, constituting between 30 and 50 square centimeters, on average, of sexually sensitive tissue according to available estimates.(54,55) That said, both the clitoral and penile prepuces serve similar functions in protecting the soft, sensitive, and moist glans tissue, preventing abrasion and drying out. In that specific regard, then, there would be a comparable effect of removing them.

“Category 3 contains those procedures that are likely to impair the ability of the recipient to engage in or enjoy sexual relations. Clitoridectomy, whether partial or complete, falls into this category.”

It is unclear whether excision of the external clitoris is “likely to impair the ability of the recipient to engage in or enjoy sexual relations,” and thus whether clitoridectomy is an apt illustration for the proposed Category 3. This is because the clitoris is a “multiplanar” organ that is nearly as long as the vagina is deep, that is, about 5-7.5 cm, with approximately 80% of its length and most of its erectile tissue being subcutaneous.(56,57) This tissue may be able to be stimulated through the vagina; it can also be stimulated externally even if the external glans has been resected, by applying pressure to the remaining tissue. There is certainly a *risk* of diminished sexual experience with clitoral excision, but as Catania et al. state: “our results suggest that FGM/C [including clitoridectomy] does not necessarily have a negative impact on psychosexual life (fantasies, desire, pleasure, ability to experience orgasm). [Even] in infibulated women, some fundamental structures for the orgasm have not been excised.”(58)

What this suggests is that it is possible to remove even a great deal of tissue from the external female genitalia and yet “leave enough behind” that there is nevertheless a decent chance that the person will be able to “enjoy sex” (as measured broadly by these kinds of studies), “experience pleasure during sexual intercourse,” and even orgasm. However, that those should be the benchmarks for acceptability is doubtful: even if it is physiologically possible to have an orgasm after one’s external clitoral glans has been excised (or to experience at least some degree of pleasure during sex due to the stimulation of other parts of the vulva/vagina that have not been removed), this does not mean that sex would be no different if one still had one’s glans. Some women who have had parts of their genitals removed in childhood—even if they can still “enjoy sex”—feel upset, angry, violated, and mutilated, simply because of the fact that part of their genitals were removed without their permission.(34) Other women who have undergone such procedures do not feel this way.(59,60) However, there is a crucial difference between these two cases. Anyone who would like to have her clitoral glans, clitoral hood, or labia removed or altered (but hasn’t yet had this done) can always undertake the surgery later; whereas, someone who did have those things done to her—but wishes they hadn’t been—has no recourse.(61) Similarly with respect to male circumcision/MGA: many men—precise numbers are hard to come by, but a recent YouGov survey concluded that fully 10% of American men wish that they

had not been circumcised(62)—have impaired sexual lives either because their surgery was “botched,” or because they feel angry, violated, mutilated, etc., in virtue of having had a large sleeve of functional, erogenous tissue removed from their genitals before they could object.(63,64)

“Critics of FGA have pointed out that there is no medical benefit to factor in the risk versus benefit calculus so often used in medicine and when compared to male circumcision. However, up to recently, the medical benefits of male circumcision were also thought to be tenuous, contested, or so minor that circumcision was classified as an elective, cosmetic procedure.”

Again, A&J are conflating adult (and hence voluntary), medicalized circumcision and infant (and hence involuntary), ritual circumcision. There is little in the way of robust data on “health benefits” as concerns the latter, and most of the data that do exist are in fact extremely “tenuous” and “contested.” In fact, the only medical bodies that have suggested that the known benefits of circumcision outweigh the known risks (though note that these bodies, too, conflate adult and infant circumcision in their assessments of the available data) are the ones mentioned by A&J in their paper. But they are in the minority on a global scale (see earlier discussion).

“The Jacobs’s three-pronged test has been previously proposed that, if satisfied, would morally preclude a government or regulatory agency from reversing a parental decision to involve a child in a minority group practice. First, the practice in question must not significantly burden either society or its members outside the group. Second, the practice must not (a) create burdens that a reasonable person outside the group would not accept for himself, or that a reasonable parent would not accept for her child (such as child marriage or slavery); or (b) carry a substantial chance of death or of major disruption of a physiological function. Third, the burden on society or individuals must be actual and substantial, not hypothetical or unlikely. ... Categories 1 and 2 of FGA ... fulfill these criteria and thus, a government or regulatory agency does not have a medical basis for interfering with a parental decision to practice a cultural or religious belief.”

First, the “Jacobs’s three-pronged test” was proposed by the same Jacobs who now advocates, with Arora, for “minor” forms of FGA, in a paper in the *Israeli Medical Association Journal* whose overriding purpose was to mount a defense of ritual male circumcision. There is no evidence that ethicists, moral philosophers, or political theorists accept this test as having any

validity, and what attention it has received has not necessarily been favorable.(65) But let us just take it for granted. Criterion #2 is that “the practice in question must not ... create burdens that a reasonable parent [outside the group] would not accept for her child.” So one can ask: Would a reasonable parent in a Western country, that is not a member of an immigrant group that practices FGA, accept for their daughter a non-therapeutic cutting procedure to the child’s vulva that carries the risk (however slight) of damaging erogenous tissue? My guess is “no,” but this is admittedly an empirical question. Finally, please note that A&J once again shift the terms of their analysis halfway through the paragraph. They begin by stating that the three-pronged test, if satisfied, would “morally preclude” a government from reversing a parental decision. But they finish by stating that a government “does not have a medical basis” for interfering with a parental decision. This careering back and forth between medical and moral arguments does not give the impression of a principled stand on the issue, but rather a desire to defend the permissibility of culturally-motivated cutting of children’s genitals (of whatever sex) by whatever argumentative means.

“Procedures that compromise sexual function, sexual enjoyment and reproductive capacity clearly violate the best interest of the child. ‘De minimis’ procedures such as removal of the clitoral hood or a ritual nick on the external female genitalia .. cause little or no functional harm.”

It seems clear that removal of the clitoral hood would cause functional harm: specifically, it would eliminate all the functions performed by the clitoral hood itself, including protection of the clitoral glans (much as the penile prepuce serves to protect the penile glans). Notice also the phrasing “little or no” harm – is a “little” harm acceptable then? And who gets to decide how much harm is only “a little”? Harm is a subjective matter. Given that A&J cannot imagine how removing something as large, sensitive, and functional as the labia minora could have a meaningfully adverse effect on a woman’s sexuality (see above discussion), it seems unlikely that their subjective judgments about sexual harm would be particularly widely shared by others. An alternative proposal is that it is the individual herself (who will be affected by the genital procedure) who should be able to decide how much value she places on having an intact vulva, and how “harmful” it would be to her sexual experience to have parts of it surgically altered or removed.

“The best interests of a child encompass not only physical well-being, but social, economic,

psychological and spiritual well-being.”

If the best interests of the child encompass not just physical well-being (which is clearly the case) but also social, economic, psychological, and spiritual well-being, then preservation of the child’s future autonomy (i.e., the freedom to make an important decision about a private part of one’s body later on) should arguably be counted among those non-physical interests that need to be factored into the equation.(66)

“The tolerant attitude in the United States to male circumcision is in stark contrast to its treatment of FGA. Yet, ‘both are likely voluntary choices influenced by cultural conditioning.’”

If either male circumcision or FGA is performed on a child before that child can understand what is at stake in the procedure, then the procedure cannot meaningfully be described as “voluntary.” It is true, however, that parents who perform (or authorize) such procedures are typically influenced by cultural conditioning.

“Neonatal boys are certainly just as vulnerable as girls. In fact, one could argue that the pubescent or adolescent girl undergoing FGA is more capable of assenting to the procedure and claiming the culture/religion as her own, than the neonatal boy. We do not condone the forcible practice of FGA if a child developmentally capable of providing assent declines to do so.”

One cannot help but agree. Neonatal boys (and girls) are uniquely vulnerable: they cannot effectively resist having a non-therapeutic procedure carried out on their genitals. But if A&J do not think that a child who is capable of declining to assent to FGA (and presumably MGA) should have the procedure forced on her (or him), then why do they approve of forcing the same procedure on an infant who is not yet *capable* of declining to assent (at least not articulately, using words)? Presumably, they would respond: “because the vulnerable infant is entirely in the parents’ care, and must depend on them to make decisions that they judge to be in its best interest.” But one could equally recommend that irreversible surgeries whose very status as being in the child’s best interest is contentious should be deferred until such a time as the individual who will actually be affected by them is in a position to weigh the pros and cons in light of her (or his) own values.(34,67)

References for the Appendix

1. Earp BD. In defence of genital autonomy for children. *J Med Ethics*. 2016;in press.
2. Arora K, Jacobs A. Female genital alteration: a compromise solution. *J Med Ethics*. 2016;in press.
3. Earp BD. Female genital mutilation and male circumcision: toward an autonomy-based ethical framework. *Medicolegal and Bioethics*. 2015;5:89–104.
4. Earp BD. Do the benefits of male circumcision outweigh the risks? A critique of the proposed CDC guidelines. *Front Pediatr*. 2015;3:18.
5. Darby R. Risks, benefits, complications and harms: neglected factors in the current debate on non-therapeutic circumcision. *Kennedy Inst Ethics J*. 2015;25(1):1–34.
6. Davis D. Male and female genital alteration: A collision course with the law? *Health Matrix*. 2001;11:487–570.
7. Svoboda JS, Darby R. A rose by any other name? Symmetry and asymmetry in male and female genital cutting. In: Zabus C, editor. *Fearful Symmetries: Essays and Testimonies Around Excision and Circumcision*. Amsterdam and New York: Rodopi; 2008. p. 251–302.
8. Dekkers W, Hoffer C, Wils JP. Bodily integrity and male and female circumcision. *Med Health Care Philos*. 2005;8(2):179–91.
9. Caldwell JC, Orubuloye IO, Caldwell P. Male and female circumcision in Africa from a regional to a specific Nigerian examination. *Soc Sci Med*. 1997;44(8):1181–93.
10. Svoboda JS. Promoting genital autonomy by exploring commonalities between male, female, intersex, and cosmetic female genital cutting. *Global Discourse*. 2013;3(2):237–55.
11. Lightfoot-Klein H. Similarities in attitudes and misconceptions about male and female sexual mutilations. In: Denniston G, Milos M, editors. *Sexual Mutilations*. Boston, MA: Springer US; 1997. p. 131–5.
12. Lightfoot-Klein H, Chase C, Hammond T, Goldman R. Genital surgeries on children before an age of consent. In: Szuchman LT, Muscarella F, editors. *Psychological Perspectives on Human Sexuality*. New York: John Wiley & Sons; 2000. p. 440–79.
13. Hellsten SK. Rationalising circumcision: from tradition to fashion, from public health to individual freedom--critical notes on cultural persistence of the practice of genital mutilation. *J Med Ethics*. 2004;30(3):248–53.
14. Gunning I. Women and traditional practices: female genital surgery. In: Askin KD, Koenig DM, editors. *Women and International Human Rights Law*. New York: Transnational Publishers; 1999. p. 651–82.
15. Ahmadu F. Rites and wrongs: An insider/outsider reflects on power and excision. In: Shell-Duncan B, Hernlund Y, editors. *Female "Circumcision" in Africa: Culture, Controversy, and Change*. Boulder, CO: Lynne Rienner Publishers; 2000. p. 283–312.

16. Fox M, Thomson M. Foreskin is a feminist issue. *Australian Feminist Studies*. 2009;24(60):195–210.
17. Kimmell MS. The kindest un-cut: emminism, Judaism, and my son’s oreskin. *Tikkun*. 2001;16(3). Available from: <http://www.cirp.org/pages/cultural/kimmell/>
18. Brusa M, Barilan YM. Cultural circumcision in EU public hospitals--an ethical discussion. *Bioethics*. 2009;23(8):470–82.
19. Earp BD. On the supposed distinction between culture and religion: A brief comment on Sir James Munby’s decision in the matter of B and G (children). *Practical Ethics*. 2015. Available from: <http://blog.practicaethics.ox.ac.uk/2015/02/on-the-supposed-distinction-between-culture-and-religion-a-comment-on-sir-james-munbys-decision-in-the-matter-of-b-and-g-children/>
20. Dustin M. Female genital mutilation/cutting in the UK: challenging the inconsistencies. *European Journal of Women’s Studies*. 2010;17(1):7–23.
21. WHO | Voluntary medical male circumcision for HIV prevention. [cited 2015 Nov 25]. Available from: http://www.who.int/hiv/topics/malecircumcision/fact_sheet/en/
22. Frisch M, Aigrain Y, Barauskas V, Bjarnason R, Boddy SA, Czauderna P, et al. Cultural bias in the AAP’s 2012 Technical Report and Policy Statement on male circumcision. *Pediatrics*. 2013;131(4):796–800.
23. Canadian Paediatric Society. Newborn male circumcision. *Paediatrics & Child Health*. 2015;20(6):311–5.
24. Earp BD. Strengths and weaknesses in the 2015 Canadian Paediatric Society statement regarding newborn male circumcision. *Paediatrics & Child Health*. 2015;in press.
25. Hartmann W. Expert statement: Dr med. Wolfram Hartmann, President of “Berufsverband der Kinder- und Jugendärzte” for the hearing on the 26th of November 2012 concerning the drafting of a federal government bill. *Berufsverband der Kinder- und Jugendärzte (BVKJ)*. 2012; Available from: http://www.kinderaerzte-im-netz.de/bvkj/kinpopup/psfile/pdf/70/121126_Ste50aa5e211e6a6.pdf
26. Royal Australasian College of Physicians. Circumcision of infant males. *Paediatrics & Child Health Division, The Royal Australasian College of Physicians*. 2010; Available from: <http://www.racp.edu.au>
27. Royal Dutch Medical Association. *Non-therapeutic circumcision of male minors*. KNMG. 2010; Available from: <http://www.knmg.nl/Publicaties/KNMGpublicatie/77942/Nontherapeutic-circumcision-of-male-minors-2010.htm>
28. Lindboe A, Malmberg F, Aula MK, Larsen P, Sigurðardóttir MM, Larsen AC, et al. Let the boys decide on circumcision: Joint statement from the Nordic Ombudsmen for Children and pediatric experts. 2013; Available from: <http://barneombudet.no/wp-content/uploads/2013/11/English-statement-.pdf>

29. National Health Service. Circumcision. NHS Choices. 2014. Available from: <http://www.nhs.uk/Conditions/circumcision/Pages/Introduction.aspx>
30. The American College of Obstetricians and Gynecologists. Newborn circumcision. 2012; Available from: <http://www.acog.org/-/media/For-Patients/faq039.pdf?dmc=1&ts=20151125T0134142988>
31. Gele AA, Bø BP, Sundby J. Have we made progress in Somalia after 30 years of interventions? Attitudes toward female circumcision among people in the Hargeisa district. *BMC Res Notes*. 2013;6:122.
32. Johnsdotter S, Essén B. Cultural change after migration: circumcision of girls in Western migrant communities. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2015;in press.
33. Leye E, Deblonde J. Legislation in Europe regarding female genital mutilation and the implementation of the law in Belgium, France, Spain, Sweden, and the UK. International Centre for Reproductive Health, Ghent University. 2004; Available from: https://www.google.co.uk/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0ahUKEwjKp431_arJAhVFWhoKHXXdAyAQFggjMAA&url=http%3A%2F%2Fec.europa.eu%2Fjustice%2Fgrants%2Fresults%2Fdaphne-toolkit%2Fen%2Ffile%2F1134%2Fdownload%3Ftoken%3DBdK4Nk5g&usg=AFQjCNFpieiDV7mWs7V9epC5IjIDyIpw8Q&sig2=6jkFhhnZmHd1XpH0ZJkz4w
34. Earp BD. Between moral relativism and moral hypocrisy: Reframing the debate on “FGM.” *Kennedy Inst. Ethics J*. 2016;in press. Available from: https://www.academia.edu/10197867/Between_moral_relativism_and_moral_hypocrisy_Reframing_the_debate_on_FGM_
35. Belluck P. Group backs ritual “nick” as female circumcision option. *The New York Times*. 2010. Available from: http://www.nytimes.com/2010/05/07/health/policy/07cuts.html?_r=1
36. Rahman A, Toubia N. *Female genital mutilation: A practical guide to worldwide laws and policies*. London: Zed Books; 2000.
37. Wald R, Braver Moss L. *Celebrating Brit Shalom*. Oakland: Notim Press; 2015.
38. Goldman R. *Questioning Circumcision: A Jewish Perspective*. Lakewood: Vanguard Publications; 1997.
39. Hernlund Y. Cutting without ritual and ritual without cutting: female “circumcision” and the e-ritualization of initiation in Gambia. In: *Female “Circumcision” in Africa: Culture, Controversy, and Change*. Boulder: Lynne Rienner Publishers; 2000.
40. Bellieni CV, Alagna MG, Buonocore G. Analgesia for infants’ circumcision. *Ital J Pediatr*. 2013 Jun 13;39:38.
41. Earp BD. Hymen “restoration” in cultures of oppression: how can physicians promote individual patient welfare without becoming complicit in the perpetuation of unjust social norms? *J Med Ethics*. 2014;40(6):431.

42. Gollaher DL. *Circumcision: A History Of The World's Most Controversial Surgery*. New York: Basic Books; 2001.
43. Stallings R, Karugendo E. Female Circumcision and HIV Infection in Tanzania: for Better or for Worse? Paper presented at the Third International AIDS Society Conference on HIV Pathogenesis and Treatment. 2005.
44. Demaria J, Abdulla A, Pemberton J, Raees A, Braga LH. Are physicians performing neonatal circumcisions well-trained? *Can Urol Assoc J*. 2013 Aug;7(7-8):260–4.
45. Cold CJ, Taylor JR. The prepuce. *British Journal of Urology: International*. 1999;83:33–44.
46. Puppo V. Anatomy and physiology of the clitoris, vestibular bulbs, and labia minora with a review of the female orgasm and the prevention of female sexual dysfunction. *Clin Anat*. 2013;26(1):134–52.
47. Schober J, Cooney T, Pfaff D, Mayoglou L, Martin-Alguacil N. Innervation of the labia minora of prepubertal girls. *J Pediatr Adolesc Gynecol*. 2010;23(6):352–7.
48. Brzyski R, Knudtson J. Female external genital organs. *Merck Manual*. 2015. Available from: <http://www.msdmanuals.com/en-gb/home/women-s-health-issues/biology-of-the-female-reproductive-system/female-external-genital-organs>
49. Jacobs AJ, Arora KS. Ritual male infant circumcision and human rights. *Am J Bioeth*. 2015;15(2):30–9.
50. Earp BD. Sex and circumcision. *Am J Bioeth*. 2015;15(2):43–5.
51. Jacobs AJ, Arora KS. Response to open peer commentaries on “ritual male infant circumcision and human rights”. *Am J Bioeth*. 2015;15(3):W1–W4.
52. Phang CW, Hocking J, Fairley CK, Bradshaw C, Hayes P, Chen MY. More than just anal sex: the potential for sexually transmitted infection transmission among men visiting sex-on-premises venues. *Sex Transm Infect*. 2008;84(3):217–9.
53. Bossio JA, Pukall CF, Bartley K. You either have it or you don't: The impact of male circumcision status on sexual partners. *Can J Hum Sex*. 2015;24(2):104–19.
54. Werker PM, Terng AS, Kon M. The prepuce free flap: dissection feasibility study and clinical application of a super-thin new flap. *Plast Reconstr Surg*. 1998;102(4):1075–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9734426>
55. Kigozi G, Wawer M, Ssettuba A, Kagaayi J, Nalugoda F, Watya S, et al. Foreskin surface area and HIV acquisition in Rakai, Uganda (size matters). *AIDS*. 2009;23(16):2209–13.
56. O'Connell HE, Sanjeevan KV, Hutson JM. Anatomy of the clitoris. *J Urol*. 2005 Oct 1;174(4 Pt 1):1189–95.
57. Baskin LS, Erol A, Li YW, Liu WH, Kurzrock E, Cunha GR. Anatomical studies of the human clitoris. *J Urol*. 1999;162(3 Pt 2):1015–20.

58. Catania L, Abdulcadir O, Puppo V, Verde JB, Abdulcadir J, Abdulcadir D. Pleasure and orgasm in women with Female Genital Mutilation/Cutting (FGM/C). *J Sex Med*. 2007;4(6):1666–78.
59. Seven things to know about female genital surgeries in Africa. *Hastings Cent Rep*. 2012;42(6):19–27. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/hast.81/full>
60. Johnsdotter S. Discourses on sexual pleasure after genital modifications: the fallacy of genital determinism (a response to J. Steven Svoboda). *Global Discourse*. 2013;3(2):256–65.
61. Carmack A, Notini L, Earp BD. Should surgery for hypospadias be performed before an age of consent? *J Sex Res*. 2015; in press.
62. Moore P. YouGov | Young Americans less supportive of circumcision at birth. YouGov. Available from: <https://today.yougov.com/news/2015/02/03/younger-americans-circumcision/>
63. Watson L. *Unspeakable Mutilations: Circumcised Men Speak Out*. New Zealand: Amazon Media; 2014.
64. Bossio J. Examining sexual correlates of neonatal male circumcision. Doctoral dissertation. 2015. Available from: <https://qspace.library.queensu.ca/handle/1974/13627>
65. Lyons B, O’Dwyer RH. The Jacobs Parental Prerogative Test. *Am J Bioeth*. 2015;15(2):52–3.
66. Darby RJ. The child’s right to an open future: is the principle applicable to non-therapeutic circumcision? *J Med Ethics*. 2013;39(7):463–8.
67. Maslen H, Earp BD, Cohen Kadosh R, Savulescu J. Brain stimulation for treatment and enhancement in children: an ethical analysis. *Front Hum Neurosci*. 2014;8:953.

Permit female genital ‘nicks’ that respect culture/religion but don’t harm, say experts

Compromise needed to protect young women from more serious forms of genital cutting

A small surgical ‘nick’ or minimalist procedures that slightly change the look, but not the function or sensory capacity of a young woman’s external genitalia, should be legally permitted as a compromise solution to the vexed issue of FGM, argue gynaecologists in the ***Journal of Medical Ethics***.

This more nuanced approach would uphold cultural and religious traditions without sacrificing the health and wellbeing of girls and young women, contend the US authors.

Despite 30 years of campaigning, the practice of cutting women’s genitalia continues to flourish in many African countries and in immigrant African communities elsewhere. To date, attempts to stamp it out with legislation have failed, and may instead be driving it underground, they suggest.

“We are *not* arguing that *any* procedure on the female genitalia is desirable,” they emphasise. “Rather, we only argue that certain procedures ought to be tolerated by liberal societies.”

To begin with, the term ‘female genital mutilation’ (FGM) should be replaced with the less emotive ‘female genital alteration’ (FGA) to reflect the different types of procedure and their associated risks, and to minimise ‘demonisation’ of important cultural practices, they say.

FGM is not an appropriate term to use for the type of procedures they advocate, which are akin to cosmetic dentistry (orthodontics), breast implants, or the type of vaginal lip sculpting (labiaplasty) “for which affluent women pay thousands of dollars,” they insist.

Current categorisation covers four types of female genital cutting, with type IV the most invasive and dangerous. But the authors call for a new system of categorisation that is based on the effects of the procedure rather than the process.

Category 1 would include procedures that should have no long lasting effects on the appearance or function of the genitalia, if performed properly: an example would be a small nick in the vulvar skin.

Category 2 would include procedures that change the appearance slightly but which are not expected to have any lasting effects on reproductive capacity or sexual fulfilment. Examples include pulling back the hood of the clitoris and labiaplasty.

Categories 3-5 would include procedures, such as clitoris removal and vaginal cauterisation that maim or harm and impair sexual fulfilment, pregnancy and childbirth. These should be banned, they say.

Categories 1 and 2 are no different to male circumcision, which is rarely performed for therapeutic benefit, but which is tolerated and legal in liberal societies, the authors argue.

And restricting these categories of FGA is “culturally insensitive and supremacist and discriminatory towards women,” they contend.

Rather, permitting this compromise would better protect girls and young women from the long term harms of the more severe forms of female genital cutting, they suggest.

“In order to better protect female children from the long term harms of categories 3 and 4 of FGA, we must adopt a more nuanced position that acknowledges that categories 1 and 2 are different in that they are not associated with long term medical risks, are culturally sensitive, do not discriminate on the basis of gender and do not violate human rights,” they conclude.

But in one of a series of commentaries in response to this paper, Professor Ruth Macklin of Albert Einstein College of Medicine, New York, insists that there is no parity between categories 1 and 2 FGA and male circumcision.

“That may be true regarding the degree of harm the procedure causes, but it is not true of the origins or the continued symbolic meaning of FGA as a necessity for being an ‘acceptable woman’,” she explains. “There is no doubt that in whatever form, FGA has its origin and purpose in controlling women.”

And she concludes: “Cultural change proceeds slowly. But with strong support from non-governmental organisations, especially those comprising local and regional women, a cultural tradition designed to control women—even in its least harmful form—is best abandoned.”

In another commentary, Brian D Earp, visiting scholar at the Hastings Center, Bioethics Research Institute in New York, argues that permitting minimalist FGA would generate a litany of legal, regulatory, medical, and sexual problems, leading to “a fiasco.”

Rather than continuing to tolerate male circumcision, and using this as a benchmark for allowing ‘minor’ forms of FGA, it may instead be time to consider taking a less tolerant stance towards both procedures, he says.

“Ultimately, I suggest that children of whatever sex or gender should be free from having healthy parts of their most intimate sexual organs either damaged or removed, before they can understand what is at stake in such an intervention and agree to it themselves,” he writes.

In a further commentary, Dr Arianne Shahvisi, of the Department of Ethics at the University of Sussex, says that a minimalist approach to FGA is unlikely to fulfil the intentions of the procedure—to change the aesthetic appearance of the female genitalia, and to control women’s sexual appetites.

And she wonders why the authors don’t take the opportunity to recommend a more minimalist approach to male circumcision.

“Rites of passage are important to all of us, but one must not cause irreversible changes to the body of another person without their consent,” she writes.

Finally, in a linked editorial, Dr Michael Dunn, of the Ethox Centre, University of Oxford, points out “The main argument is controversial, but its airing on the pages of the journal has a clear purpose: by subjecting FGM in its many forms to ethical analysis, we will be in a stronger position to develop and tailor interventions that function to prevent indefensible practices of this kind.”

The evidence suggests that at least 200 million girls and women alive today have been subjected to genital cutting, he says.